

## **UC Irvine**

### **Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health**

#### **Title**

Olivier Syndrome: Traumatic Asphyxia

#### **Permalink**

<https://escholarship.org/uc/item/06g0b348>

#### **Journal**

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 14(6)

#### **ISSN**

1936-900X

#### **Authors**

Shiber, Joseph R  
Fontane, Emily  
Monroe, Brett

#### **Publication Date**

2013

#### **DOI**

10.5811/westjem.2013.7.18581

#### **Copyright Information**

Copyright 2013 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

## Olivier Syndrome: Traumatic Asphyxia

Joseph R. Shiber, MD  
Emily Fontane, MD  
Brett Monroe, MD

University of Florida, College of Medicine Jacksonville, Department of Emergency  
Medicine, Jacksonville, Florida

*Supervising Section Editor:* Sean Henderson, MD

Submission history: Submitted June 16, 2013; Accepted July 1, 2013

Electronically Published October 17, 2013

Full text available through open access at [http://escholarship.org/uc/uciem\\_westjem](http://escholarship.org/uc/uciem_westjem)

DOI: 10.5811/westjem.2013.7.18581

[West J Emerg Med. 2013;14(6):653.]

A 51 year-old woman was found confused while crawling across a field. She had marked facial cyanosis and edema with cutaneous petechiae, subconjunctival hemorrhages, and ecchymosis across her anterior neck (image 1). Mild cerebral edema and a non-displaced thyroid cartilage fracture were found on computed tomography (CT). The patient recovered full neuro-cognitive function within 24 hours and reported that she had been assaulted and choked by the throat. Her airway remained stable and the laryngeal injury was treated conservatively; she was discharged home after three days.

Olivier described this syndrome over 150 years ago after thoraco-abdominal crush injuries. It is essentially a prolonged valsalva maneuver that results in increased venous pressure and stasis above the level of the compressive force.<sup>1,2</sup> The consequent findings on the chest, neck and face are startling but of no prognostic significance, and the majority of patients have a favorable outcome.<sup>2,3</sup>

---

*Address for Correspondence:* Joseph R Shiber, MD, University of Florida, College of Medicine Jacksonville, 1st Floor, Clinical Center, 655 West 8th Street, Jacksonville, FL 32209. Email: joseph.shiber@jax.ufl.edu.

---

*Conflicts of Interest:* By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.



**Figure.** Patient with facial cyanosis and edema with cutaneous petechiae, subconjunctival hemorrhages, and ecchymosis across the anterior neck.

### REFERENCES

1. Jongewaard W, Cogbill T, Landercasper J. Neurologic consequences of traumatic asphyxia. *J Trauma*. 1992;32(1):28-31.
2. Sarihan H, Abes M, Akyazici R, et al. Traumatic asphyxia in children. *J Cardiovasc Surg*. 1997;38(1):93-5.
3. Richards C, Wallis D. Asphyxiation: a review. *Trauma*. 2005;7:37-45.