

# UCLA

## UCLA Previously Published Works

### Title

Reply Paradoxical Brain Embolism, Not Only From a Patent Foramen Ovale

### Permalink

<https://escholarship.org/uc/item/23603414>

### Journal

JACC Cardiovascular Interventions, 10(3)

### ISSN

1936-8798

### Authors

Kijima, Yasufumi

Rafique, Asim M

Tobis, Jonathan M

### Publication Date

2017-02-01

### DOI

10.1016/j.jcin.2016.12.001

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## Letters

### TO THE EDITOR

## Paradoxical Brain Embolism, Not Only From a Patent Foramen Ovale



We read with interest an article by Kijima et al. (1), which highlights the fact that not only a patent foramen ovale (PFO) can be responsible for a cryptogenic stroke. Right-to-left shunting (RLS) through an PFO is undeniably the major cause of paradoxical embolism. However, rarely, it can also be caused by RLS at pulmonary level due to pulmonary arteriovenous malformation (PAVM). A PAVM is a direct communication between the branches of the pulmonary artery and pulmonary veins. Large or multiple PAVMs can cause cyanosis and exertional dyspnea, but usually patients are asymptomatic. Regardless of their size, PAVMs can affect the central nervous system causing migraines, transient ischemic attacks, strokes, or abscess. The reported incidence of strokes in patients with PAVM is 18% to 32% and up to 60% in cases of multiple PAVMs (2). It has also been shown that recurrent strokes occur more often in patients with PAVMs than with PFO (probably because of continuous RLS). Classical diagnostic tools used to confirm PAVM are contrast-enhanced computed tomography or magnetic resonance imaging and pulmonary angiography. Chest radiography suggests PAVM only in about 45% of patients.

It should be underscored that PAVMs can be also diagnosed using contrast transesophageal echocardiography (c-TEE) and contrast transcranial Doppler (c-TCD), which are considered the “gold standard” for revealing an PFO. In the case of an extracardiac shunt, c-TEE with a Valsalva maneuver shows bubbles entering the left atrium 3 to 8 cardiac cycles after they were seen in the right atrium. In contrast, in cardiac RLS, the “3-beat rule” is used, which means that bubbles should appear in the left atrium between first and third cardiac cycles (3). In addition, RLS can be identified by the use of c-TCD. The technique is based on the detection of an intravenously injected contrast within intracranial arteries. In case of an RLS, the contrast enters the arterial circulation and produces microembolic signals. Microembolic signals

passing pulmonary shunts appear later in the cerebral circulation than those passing cardiac shunts. The time window characteristic for PAVMs is about 15 s (11 s for intracardiac shunts), but it depends on the heart rate (duration of about 6 heart beats) (3).

In conclusion, it is of great importance to distinguish the level of RLS in c-TEE and c-TCD. It seems that PAVM can be responsible for some “false-positive” results of c-TEE or c-TCD. Finally, it is reasonable to consider c-TCD or c-TEE after every PFO closure to identify potential persistent RLS. For complete prevention of recurrent strokes caused by paradoxical embolism, it is necessary to not only close a PFO, but all existing shunts.

Robert Sabiniewicz, MD

\*Lidia Wozniak-Mielczarek, MD

\*Department of Pediatric Cardiology and Congenital Heart Diseases

Medical University Gdansk, Poland

7 Debinki Street

Gdansk 80-211

Poland

E-mail: [lidiawozniak@yahoo.com](mailto:lidiawozniak@yahoo.com)

<http://dx.doi.org/10.1016/j.jcin.2016.11.051>

Please note: The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

### REFERENCES

1. Kijima Y, Rafique AM, Tobis JM. Patent foramen ovale combined with pulmonary arteriovenous malformation. *J Am Coll Cardiol Intv* 2016;9: 2169-71.
2. White RI Jr., Lynch-Nyhan A, Terry P, et al. Pulmonary arteriovenous malformations: techniques and long-term outcome of embolotherapy. *Radiology* 1988;169:663-9.
3. Jauss M, Zanette EM. Detection of right-to-left shunt with ultrasound contrast agent and transcranial Doppler sonography. *Cerebrovasc Dis* 2000; 10:490-6.

### REPLY: Paradoxical Brain Embolism, Not Only From a Patent Foramen Ovale



We appreciate the valuable commentary on our paper by Drs. Sabiniewicz and Wozniak-Mielczarek. A right-to-left shunt (RLS) caused by a pulmonary arteriovenous malformation (PAVM) and that caused by a patent foramen ovale (PFO) is most often diagnosed with echocardiographic modalities using agitated saline bubble contrast techniques. These include contrast transthoracic echocardiography, contrast transesophageal echocardiography, and contrast

transcranial Doppler (TCD). For contrast transthoracic echocardiography and transesophageal echocardiography, a PFO is distinguished from bubbles passing through the pulmonary circulation based on the timing of the appearance of contrast in the left atrium, which is called the “3-beat rule.” However, there are exceptions to this general rule. A large PAVM can produce early appearance of bubbles that look exactly like what is seen with a PFO (1). Therefore, the rules defined by the timing of the appearance of bubbles tend to be misleading and often provide an inadequate diagnosis. Theoretically, when a PFO is combined with a PAVM, bubbles through a PFO appear in the left atrium before the appearance of bubbles through a PAVM.

TCD can be performed with an adequate Valsalva maneuver especially when aided by visual feedback using a manometer, and is the most sensitive method to detect an RLS (2). One criticism of TCD is that it does not distinguish the etiology of an RLS. However, the degree of an RLS caused by a PFO tends to increase from rest to Valsalva, whereas an RLS caused by a PAVM does not (3). This is another reason why quantitative measurement of the degree of an RLS with TCD is preferable to transesophageal echocardiography.

Distinguishing the level of an RLS is important and a PAVM can mimic an RLS caused by a PFO. It is not straightforward to distinguish one from the other and sometimes (20%) these 2 etiologies coexist. A PFO or PAVM should be closed as secondary prevention for cryptogenic stroke, but either should be closed as primary treatment for hypoxemia.

Yasufumi Kijima, MD

Asim M. Rafique, MD

\*Jonathan M. Tobis, MD

\*Program in Interventional Cardiology

Department of Medicine

David Geffen School of Medicine at UCLA

10833 Le Conte Avenue

Factor Building B-976

Los Angeles, California 90095

E-mail: [jtobis@mednet.ucla.edu](mailto:jtobis@mednet.ucla.edu)

<http://dx.doi.org/10.1016/j.jcin.2016.12.001>

Please note: Dr. Tobis serves as a consultant for W.L. Gore and Associates, Inc., and St. Jude Medical, Inc. The other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

## REFERENCES

1. Homma S, Messé SR, Rundek T, et al. Patent foramen ovale. *Nat Rev Dis Primers* 2016;2:15086.
2. Van H, Poommipanit P, Shalaby M, Gevorgyan R, Tseng CH, Tobis J. Sensitivity of transcranial Doppler versus intracardiac echocardiography in the detection of right-to-left shunt. *J Am Coll Cardiol Img* 2010;3:343-8.
3. Kijima Y, Gevorgyan R, McWilliams JP, Miller N, Noureddin N, Tobis JM. Usefulness of transcranial Doppler for detecting pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia. *Am J Cardiol* 2016;117:1180-4.