

UC Agriculture & Natural Resources

California Agriculture

Title

Using Ecological Site Descriptions to make ranch-level decisions about where to manage for soil organic carbon

Permalink

<https://escholarship.org/uc/item/3218t061>

Journal

California Agriculture, 76(2-3)

ISSN

0008-0845

Authors

Aoyama, Lina
Bartolome, James
Silva, Lucas
[et al.](#)

Publication Date

2022

DOI

10.3733/ca.2022a0007

Copyright Information

Copyright 2022 by the author(s). All rights reserved unless otherwise indicated. Contact the author(s) for any necessary permissions. Learn more at <https://escholarship.org/terms>

Peer reviewed

Using Ecological Site Descriptions to make ranch-level decisions about where to manage for soil organic carbon

Rangeland conservation can keep carbon out of the atmosphere by storing it in the soil. Ecological Site Descriptions can help determine promising sites.

by Lina Aoyama, James W. Bartolome, Lucas Silva and Whendee L. Silver

Online: <https://doi.org/10.3733/ca.2022a0007>

California's rangelands cover approximately 57 million acres of grasslands, shrublands, woodlands and deserts (FRAP 2018). As the most extensive land use type in California, rangelands play an important role in climate change mitigation by storing considerable amounts of soil organic carbon (SOC) below ground (Herrero et al. 2016; Sanderson et al. 2020; Schuman et al. 2002). In wildfire and drought-prone California, rangelands are considered more reliable for carbon storage because they secure most of the carbon in the soil instead of in aboveground plants that could potentially burn (Dass et al. 2018).

The primary way that carbon is stored in the soil is as soil organic matter (SOM). Soil organic matter is a mix of decomposing plant and animal biomass, soil microbes and humus. Increasing SOM also enhances soil water-holding capacity and improves nutrient cycling, which can maintain or increase forage production (Conant et al. 2001; Herrick and Wander 1997). In California, as elsewhere, there has been a growing interest in improving rangeland management practices to increase the amount of

Abstract

Maintaining and enhancing soil organic carbon storage can mitigate climate change while promoting forage growth. California has adopted incentive programs to promote rangeland practices that build soil organic carbon. However, there is no standard framework for assessing the baseline level of soil organic carbon at the ranch scale. Here, we use the Ecological Site Description — a land-type classification system — to help ranch managers set priorities about where to implement practices to increase soil organic carbon. We measured baseline carbon stocks at 0 to 15 and 15 to 30 centimeters' depth across three ecological sites and two vegetation states (shrubland and grassland) at Tejon Ranch, California. We discovered increased levels of soil carbon at ecological sites in higher elevations, and more soil carbon in shrublands as compared to grasslands. Slope, elevation, and soil texture, as well as plant litter and shrub cover, were significant predictors of soil carbon. The Ecological Site Description framework can serve as an important tool to help range managers keep carbon in the soil and out of the atmosphere.

Intersection of shrubland and grassland at the foothills of Tehachapi Mountains, Tejon Ranch, Calif. The authors found that soil carbon significantly varied by ecological sites and vegetation states.

Photo: Lina Aoyama.

Ecological Site Descriptions are a useful framework for assessing baseline soil organic carbon on diverse landscapes.

SOC (Bradford et al. 2019; Byrnes et al. 2017; Carey, Gravuer et al. 2020).

The California Department of Food and Agriculture (CDFA)'s Healthy Soils Program is one of the programs that has been put in place to promote carbon storage. This program provides financial incentives through cost sharing and technical support for land managers to implement agricultural management practices that augment SOC or prevent erosion (CDFA 2016). Between 2017 and 2020, 604 landowners participated in this program, and the estimated greenhouse gas reduction was 109,089 tonnes CO₂ eq/year (CDFA 2021). Recommended range management practices include planting oak trees, applying compost, and restoring riverbanks (Dahlgren et al. 1997; Matzek et al. 2018; Ryals et al. 2014).

Ideally, land managers should be able to easily find out the baseline amount of soil organic carbons on their lands and evaluate where to implement these practices. However, the amount of SOC varies greatly from place to place because of California's diverse climate, topography and soil conditions (Carey, Weverka et al. 2020). The Rangeland Monitoring Network is currently evaluating the variability in soil carbon storage potential across California rangelands (Carey et al. 2020b). There are SOC estimates by soil classification available on the USDA Web Soil Survey, but we lack a framework to relate these soil carbon estimates to specific landscape features at a ranch scale.

Ecological Site Description (ESD), developed and maintained by the USDA Natural Resource Conservation Service, is a land-type classification system that could facilitate ranch-level planning to manage for soil carbon. The ESD framework has primarily been used to manage vegetation on rangelands and forests, as well as to identify priorities for ecosystem benefits such as wildlife habitat, water quality and wildfire protection (Brown and Havstad 2016). This framework has not yet been extended to manage SOC stocks. An ecological site is defined as an area 24.7 to 247 acres with similar "physical properties [climate, soils, topography, underlying geologic material], potential vegetation, and responses to management that differ from other kinds of land" (USDA 2018). Each ecological site contains a state-and-transition (STM) model of vegetation states that is helpful in evaluating the effects of management actions or disturbances on the existing state of vegetation (Briske et al. 2005).

A ranch can have more than one ecological site. A related framework called Major Land Resource Area (MLRA) has been used by the National Resource

Inventory Soil Monitoring Network to estimate SOC stocks at a national scale (Ogle et al. 2010; Spencer et al. 2012). However, these estimates are not precise enough for ranch-level decision making. While similar in concept, ESD units are smaller, more site-specific than MLRAs and may be more fitting for ranch-level planning.

The goal of this study is to explore the use of ESDs for ranch-level identification of potential priority sites for soil carbon sequestration projects. We used Tejon Ranch in Kern County, California, as a case study. Tejon Ranch is an ideal place to study ESDs because it has complex biogeographical features, and its range managers already use ESDs to manage forage for livestock (Spiegel et al. 2016). Studies connecting soil carbon to plant communities are scarce in Southern California (Booker et al. 2013; Carey et al. 2020b). To examine how soil carbon content is related to ESDs and vegetation states, we measured topsoil carbon in three ecological sites at different elevations and in two dominant vegetation states. We then explored how plant and soil characteristics were related to patterns in SOC stocks at ranch level.


Ecological sites on Tejon Ranch

Tejon Ranch (270,000 acres) is the largest contiguous, privately owned property in California, located 35 miles south of Bakersfield, California (latitude 34.935044°, longitude 118.670405°). We studied the northwestern portion of the ranch (50,000 acres) where the San Joaquin Valley and Tehachapi Mountains meet. Using the ESD framework, three ecological sites were defined based on slope, elevation and geology (fig. 1): Holocene Flats, Lower Miocene Hills and Upper Miocene Hills. Tejon Ranch has been grazed by livestock since the 1840s when the original Tejon Ranchos were created (Latta 1976). During the study, beef cattle (*Bos taurus*) grazed year-round while no grazers, including wildlife, were excluded from the study area.

Dominant vegetation types in the study site were native shrubland and non-native annual grassland. Native shrub species commonly found in shrublands included cattle saltbrush (*Atriplex polycarpa*), Interior California buckwheat (*Eriogonum fasciculatum* var. *polifolium*) and Valley bladderpod (*Peritoma arborea* var. *globose*) (Aoyama et al. 2020). The grassland state was primarily composed of native and non-native annual forbs (e.g., *Plagiobothrys canescens*, *Erodium cicutarium*) and non-native annual grasses (e.g., *Bromus diandrus*, *Avena barbata*) (Spiegel et al. 2016).

Soils were generally shallow Mollisols with bedrock encountered at approximately 30 centimeters' (cm) depth. Parent materials were sedimentary rocks from the Miocene and alluvium deposits from the Holocene (Dibblee 2008). Soil classification varied from Psamments-Xerolls complex (loamy sand), to

Ecological sites



Soil sampling scheme :

Depths: 0–15 cm, 15–30 cm / Plots: 12 / Replicates: 3
Number of times sampled: 4 / $n = 288$




FIG. 1. Ecological sites along the elevational gradient of Tehachapi Mountain in Tejon Ranch, California: San Joaquin Valley Holocene Flats, Lower Miocene Hills and Upper Miocene Hills. The schematic is adapted from Spiegel et al. (2016). Each ecological site has two vegetation types: shrubland (top) and grassland (bottom). Two plots in each vegetation type are nested within each ecological site. Each plot was sampled for soil, cattle use and vegetation cover. Three replicates of soil cores were collected at each plot four times. Soils in shrubland plots were collected from three microhabitats: base of shrub, edge of shrub canopy (~0.5 m from the base) and grass patch between shrubs (~2 m from the base). Soils in grassland plots were collected randomly.

TABLE 1. Heterogeneous soil properties across the landscape

Ecological Site	pH	PO ₄ ³⁻	K ⁺	Na ⁺	Ca ²⁺	Mg ²⁺	SO ₄ ²⁻	Sand	Clay
	[H ⁺]	ppm	Meq/100 g	Meq/100 g	Meq/100 g	Meq/100 g	ppm	%	%
SJV Holocene Flats	7.56 (0.70)	5.24 (2.15)	0.66 (0.51)	0.05 (0.06)	7.96 (5.69)	1.18 (0.82)	3.53 (3.33)	83.23 (8.70)	7.24 (3.86)
Lower Miocene Hills	7.64 (0.32)	4.45 (1.48)	0.58 (0.69)	1.53 (2.99)	11.38 (7.71)	1.38 (0.39)	8.25 (12.12)	74.47 (13.26)	9.80 (4.97)
Upper Miocene Hills	7.32 (0.67)	3.52 (1.95)	0.62 (0.15)	0.24 (0.10)	23.57 (5.52)	3.50 (1.58)	5.57 (2.41)	60.82 (5.14)	20.86 (7.82)
ANOVA (<i>P</i> -value)	0.70	0.71	0.78	0.48	0.04	0.04	0.72	0.08	0.03

Values are mean and standard error of each soil property by ecological sites.

The last row contains *P*-values from ANOVAs of soil properties with ecological site as a main effect. Significant *P*-values (*P* ≤ 0.05) are bolded.

Haploxerolls (sandy loam), to Chanac-Pleito complex (sandy clay loam) (USDA Web Soil Survey); the clay content increased with elevation (table 1).

We established twelve plots (65 feet by 16 feet) stratified by three ecological sites and two dominant vegetation types (grassland versus shrubland; fig. 1). We measured plant cover (shrub, herbaceous, litter, bare and other) and cattle use (evidence of herbivory, trailing, trampling, and old and new cattle manure) in each plot in March, June and October 2017 and March 2018. In addition to qualitative measures of cattle use, we set up motion-triggered camera traps on t-posts at heights of 1.5 meters on each plot to measure the number of cattle visits from March 2017 to March 2018.


To measure soil carbon concentrations and content, we sampled three replicate cores in each plot from two depths (0–15 cm and 15–30 cm) using a 5-cm diameter by 15-cm-long soil corer. We sampled the topsoil because it is the fraction of soil that is

most accessible, where management has the most impact on soil carbon on an annual basis (Fontaine et al. 2007; Gregory et al. 2016; Syswerda et al. 2011). We sampled soils within grassland plots randomly; within shrubland plots, we stratified by microhabitats at the base of shrub main stem, at the edge of shrub canopy, and at a grass patch 2 meters from the shrub main stem. Cores were weighed fresh and after drying at 105°C for bulk density determination (see online technical appendix). We measured the concentration of SOM using the loss-on-ignition method and the concentration of SOC using the flash combustion method. We then converted concentration percentages to stocks (Mg/ha) by multiplying the values by bulk density and depth. SOC was positively correlated with SOM [$\log(\text{SOC Mg/ha}) = 0.68 * \log(\text{SOM Mg/ha}) + 0.18$; $R^2 = 0.41$, $P < 0.001$].

Soil carbon by vegetation and ecological site

Our study explored the idea of using ESDs to provide a first approximation of baseline soil carbon at ranch level. In the study area on Tejon Ranch, average SOC stocks at 0–15 cm, 15–30 cm and 0–30 cm depths were 19.18 ± 0.90 Mg C/ha, 14.25 ± 0.67 Mg C/ha and 33.63 ± 1.49 Mg C/ha, respectively (table 2). These estimates were less than the statewide average (58 ± 4 Mg C/ha in 0–25 cm) reported by Silver et al. (2010). In contrast, these values are comparable to the statewide average (19.31 Mg C/ha in 0–10 cm) reported by Carey et al. (2020) and the 20.98 Mg C/ha for the A horizon in the Sierra Nevada foothills (Eastburn et al. 2017). From the linear mixed-effect models, we found that SOM and SOC significantly varied by ecological sites and vegetation states (table 3).

Both SOM and SOC stocks increased with elevation. They were higher in the Upper Miocene Hills than in the San Joaquin Valley Holocene Flats ecological sites (table 2). This effect of elevation could be due to a combination of time, soil type, and legacy of historical plant communities. Soils in the Miocene Hills were older than those in the Holocene Flats



Mary McDonnell, Dylan Stover and Tara Harmon (from right to left) taking turns to collect soil in a grassland plot, Tejon Ranch, Calif. Photo: Lina Aoyama.

TABLE 2. Bulk density (g/cm³), soil organic matter (SOM) concentration (%), soil organic carbon (SOC) concentration (%), SOM content (Mg/ha) and SOC content (Mg/ha) by depth and spatial scale

Depth	Scale	Bulk density	SOM concentration	SOC concentration	SOM content	SOC content	
		(g/cm ³)	(%)	(%)	(Mg/ha)	(Mg/ha)	
0–15 cm	Landscape	1.11 ± 0.01	3.13 ± 0.12	1.18 ± 0.05	49.09 ± 1.64	19.18 ± 0.90	
	ES	SJV_Holo	1.17 ± 0.01	2.24 ± 0.13	0.91 ± 0.06	39.38 ± 2.19	15.81 ± 0.77
		Low_Mio	1.20 ± 0.01	2.75 ± 0.22	1.06 ± 1.13	46.73 ± 3.30	19.64 ± 2.42
		Up_Mio	0.94 ± 0.01	4.42 ± 0.13	1.58 ± 0.06	61.68 ± 2.12	22.26 ± 0.93
	Veg	Grassland	1.17 ± 0.01	2.69 ± 0.16	0.89 ± 0.05	44.49 ± 1.85	14.87 ± 0.67
		Shrubland	1.05 ± 0.01	3.54 ± 0.17	1.45 ± 0.08	53.49 ± 2.60	23.29 ± 1.53
15–30 cm	Landscape	1.13 ± 0.01	0.84 ± 0.03	0.84 ± 0.03	37.84 ± 1.17	14.25 ± 0.67	
	ES	SJV_Holo	1.20 ± 0.01	1.77 ± 0.08	0.73 ± 0.04	33.08 ± 1.39	12.87 ± 0.65
		Low_Mio	1.23 ± 0.01	1.92 ± 0.14	0.79 ± 0.08	34.08 ± 2.26	14.24 ± 1.78
		Up_Mio	0.95 ± 0.01	3.44 ± 0.13	1.03 ± 0.04	47.77 ± 1.84	15.81 ± 0.60
	Veg	Grassland	1.18 ± 0.01	2.13 ± 0.15	0.60 ± 0.03	35.95 ± 1.83	10.42 ± 0.46
		Shrubland	1.09 ± 0.01	2.51 ± 0.10	1.05 ± 0.05	39.52 ± 1.50	17.64 ± 1.07
0–30 cm	Landscape				87.44 ± 2.49	33.63 ± 1.49	
	ES	SJV_Holo			72.46 ± 2.64	28.68 ± 1.21	
		Low_Mio			81.78 ± 5.01	35.02 ± 4.02	
		Up_Mio			110.0 ± 3.39	37.66 ± 1.39	
	Veg	Grassland			80.91 ± 3.34	25.48 ± 0.99	
		Shrubland			93.46 ± 3.57	41.14 ± 2.47	

Values are mean and standard error of each estimate. ES = ecological site, Veg = vegetation state.

TABLE 3. Ecological sites and vegetation types influenced bulk density, soil organic matter (SOM) and soil organic carbon (SOC)

Response	Depth	Ecological site		Vegetation state		Ecological site: vegetation state	
		F	P-value	F	P-value	F	P-value
Bulk density	0–15 cm	8.576	0.017	4.399	0.080	0.091	0.914
	15–30 cm	6.367	0.032	1.351	0.289	0.023	0.976
SOM (Mg/ha)	0–15 cm	3.543	0.009	1.565	0.257	2.691	0.146
	15–30 cm	15.043	0.004	1.194	0.316	14.283	0.005
	0–30 cm	6.357	0.033	1.069	0.341	4.680	0.059
SOC (Mg/ha)	0–15 cm	1.324	0.033	6.386	0.044	1.949	0.222
	15–30 cm	0.541	0.607	13.154	0.011	1.465	0.303
	0–30 cm	1.744	0.252	10.393	0.018	1.8025	0.243

F statistics and P-values are results of linear mixed-effect models of SOM and SOC with ecological site, vegetation state, and their interactions as main effects and plot as a random effect. Significant P-values ($P \leq 0.05$) are bolded.

(Dibblee 2008), and thus had more time to accumulate soil carbon. High clay content found in higher elevation sites also may have contributed to higher soil carbon storage due to carbon-mineral associations (Kaiser et al. 2012). Additionally, oak savanna historically dominated the slopes of the Tehachapi Mountains (Twisselman 1967), which likely contributed to higher soil carbon inputs on the slopes than on the flats (Dahlgren et al. 1997; Koteen et al. 2011).

We also found that both SOM and SOC stocks were significantly higher in shrubland states compared to grassland states in the 0–30 cm profile (table

2), which corroborates the broad pattern found in California’s rangelands that woody plants increase soil carbon pools (Silver et al. 2010). One possible explanation for this is that shrubs in semi-arid environments tend to have dense root mats in the top 30 cm to capture transient soil moisture (Chabbi et al. 2009; Swanston et al. 2005), which contribute to soil carbon storage. Another possible explanation is that litter fall from shrubs contributes significantly to near-surface soil carbon pools (Rau et al. 2009). These results suggest that grouping the vegetation states by dominant functional groups within




FIG. 2. Soil organic carbon (SOC) stocks in 0–30 cm depth by ecological sites and microhabitats: shrub, edge of shrub canopy, grass patch between shrubs, and grassland. Horizontal bars are medians, boxes are the 25th and 75th percentile, and the whiskers are the maximum and minimum values. Dots represent outliers in the data.

ecological sites could be valuable for evaluating SOC stocks at ranch level.

Soil carbon varied across microhabitats

Given the wide range of sizes of ranches in California, the scalability of plot-level information is important. The spatial variability of SOC stocks within a ranch informs managers about the spatial scale at which they should aggregate the estimates of SOC from the USDA Web Soil Survey or their own soil sampling. The average coefficient of variance of SOC stocks in shrubland was higher (85.9) than that in grassland (38.67). This plot-level variability came from the microhabitat types (shrub base, edge of shrub canopy, grass patch between shrubs, and grassland). Estimates of SOC stocks in 0–30 cm were significantly higher at the base of shrubs than in grasslands ($F_{3,119} = 7.924, P < 0.001$; fig. 2). These results support the notion of shrubs acting as “islands of fertility” where soil carbon and nutrients are high relative to areas outside the canopy (Schlesinger et al. 1996; Schade et al. 2003; Schade and Hobbie 2005). The implication for SOC monitoring is that sampling more cores in shrubland than in grassland is necessary to capture the spatial variability in soil carbon within vegetation states.

Predictors of carbon across the landscape

Grazing management has been proposed as a means to sequester soil carbon (Byrnes et al. 2017).

However, in the literature, cattle use is often not well quantified beyond presence-absence data. We used the standard qualitative assessment of cattle use (e.g., evidence of herbivory, fresh and old cow manure, trampling) and quantified the frequency and intensity of cattle use via camera traps. We found that none of these grazing use metrics meaningfully captured variation in SOM or SOC stocks. This result is in line with prior work showing that the influence of grazing on soil carbon is not significant in semi-arid rangelands of California (Biggs and Huntsinger 2021), especially in sandier soils (Silver et al. 2010; Stanton et al. 2018).

To identify important factors that predict soil carbon at ranch level, we examined the relationships between environmental variables and soil carbon stocks. We found that mean percent cover of litter and shrubs predicted SOM stocks in the 0–30 cm profile, while elevation, slope and soil calcium content predicted SOC stocks in the 0–30 cm profile (table 4). These were similar to the factors that influenced soil carbon at the regional level (Carey et al. 2020). Managers may use this information to group their ranch or management unit by ecological sites and vegetation states, and make decisions about where on the landscape they might want to implement carbon sequestration projects. For example, managers on Tejon Ranch could use the ESDs to prioritize conserving or restoring native shrubs in the high elevation Upper Miocene Hills ecological site. On other ranches, land managers might increase soil carbon by planting oak trees in ecological sites that have oak woodland as a reference state, or by applying compost in ecological sites with low baseline

TABLE 4. Abiotic and biotic environmental factors influence soil organic matter (SOM) and soil organic carbon (SOC)

Response	Depth	Predictor	Coefficient	t-value	P-value	R ² _m	R ² _c
SOM	0–15 cm	Mg	0.458	1.732	0.121	0.489	0.963
		Litter % cover	0.514	1.675	0.132		
		Shrub % cover	0.454	1.803	0.109		
	15–30 cm	pH	0.244	1.442	0.208	0.602	0.964
		Ca	–0.399	–1.865	0.121		
		Clay	0.276	1.754	0.139		
		Slope	0.273	2.392	0.062		
Litter % cover		0.732	3.843	0.012			
Shrub % cover	0.421	3.217	0.023				
SOC	0–15 cm	Ca	0.622	0.268	0.059	0.419	0.945
		Clay	–0.557	0.304	0.116		
		Elevation	–0.989	0.361	0.034		
		Slope	0.953	0.253	0.009		
		Litter % cover	0.637	0.300	0.077		
	15–30 cm	Ca	0.528	3.306	0.013	0.396	0.990
		Clay	–0.431	–2.596	0.035		
		Elevation	–0.375	–2.084	0.075		
		Slope	0.501	3.375	0.011		

Coefficients, t-values, P-values, R²_m and R²_c are results of linear regression models with SOM and SOC as response variables, environmental factors as fixed effects, and quadrats within plots as random effects. R²_m is a proportion of variance explained by fixed effects and R²_c is that explained by both fixed and random effects. Significant P-values ($P \leq 0.05$) are bolded.

SOC stocks. This approach is compatible with range management in California that values multiple ecosystem services supported by the landscape rather than single-purpose management (Biggs and Huntsinger 2021).

Deciding where to build up carbon

California is leading the way in building up carbon in the soil, where it helps hold nutrients and water, and out of the atmosphere, where it contributes to global warming. Planting oak trees on range land, adding compost, and maintaining rivers can increase carbon sequestration while improving forage. A simple framework to assess soil organic carbon stocks would make it easier for land managers to determine where to prioritize implementing practices that increase the amount of carbon stored in the soil. In this case study at Tejon Ranch, we demonstrated that Ecological Site Descriptions are a useful framework for assessing baseline soil organic carbon on diverse landscapes. We found that the amount of soil organic carbon differs by dominant vegetation functional groups and ecological sites. Environmental factors such as soil type, topography and vegetation cover are predictors of the amount of soil carbon.

ESDs are less developed in California than in other states. However, this framework can contribute to our understanding of the environmental factors

that influence soil organic carbon. This should enable rangeland managers to focus soil management practices where they will do the most good. [CA](#)

L. Aoyama is Ph.D. student, Environmental Studies Program and Department of Biology, University of Oregon, Eugene, Oregon; J.W. Bartolome is Emeritus Professor, and W.L. Silver is Professor, Department of Environmental Science, Policy, and Management, UC Berkeley; L. Silva is Associate Professor, Environmental Studies Program, Department of Geography, and Institute of Ecology and Evolution, University of Oregon, Eugene, Oregon.

We thank Peter Hopkinson, Felix Ratcliff, Ellery Mayence, Atalie Brown, Andrew Capistrano, Jack Cronin, Denise Defreese, Lee Fram, Luke Macaulay, Mary McDonnell, Tara Harmon, Michael Orellana, Kea Rutherford, Allison Rofe, Dylan Stover and Lulu Waks for helping with fieldwork. We thank Summer Ahmed for assisting with soil analysis. Financial support was provided by the Tejon Ranch Conservancy, Range Block Grant, Russell Rustici Chair in Range Management, project numbers CA-B-ECO-0117-MS and CA-B-ECO-7673-MS of the USDA National Institute of Food and Agriculture, the University of California, Berkeley, and The Nature Conservancy under the Oren Pollak Memorial Student Research Grant for Grassland Science.

References

- Aoyama L, Bartolome JW, Hallett LM. 2020. Incorporating diversity measures into Ecological Site Descriptions to manage biodiversity on heterogeneous landscapes. *Rangelands* 42(4):93–105. <https://doi.org/10.1016/j.rala.2020.05.002>
- Biggs NB, Huntsinger L. 2021. Managed grazing on California annual rangelands in the context of state climate policy. *Rangeland Ecol Manag* 76:56–68. <https://doi.org/10.1016/j.rama.2021.01.007>
- Booker K, Huntsinger L, Bartolome JW, et al. 2013. What can ecological science tell us about opportunities for carbon sequestration on arid rangelands in the United States? *Global Environ Chang* 23:240–51. <https://doi.org/10.1016/j.gloenvcha.2012.10.001>
- Bradford MA, Carey CJ, Atwood L, et al. 2019. Soil carbon science for policy and practice. *Nat Sustain* 2:1070–2. <https://doi.org/10.1038/s41893-019-0431-y>
- Briske DD, Fuhlendorf SD, Smeins FE. 2005. State-and-transition models, thresholds, and rangeland health: A synthesis of ecological concepts and perspectives. *Rangeland Ecol Manag* 58:1–10. www.jstor.org/stable/3899791
- Brown JR, Havstad KM. 2016. Using ecological site information to improve landscape management for ecosystem services. *Rangelands* 38(6):318–21. <https://doi.org/10.1016/j.rala.2016.10.011>
- Byrnes R, Eviner V, Kebreab E, et al. 2017. Review of research to inform California's climate scoping plan: Agriculture and working lands. *Calif Agr* 71:160–8. <https://doi.org/10.3733/ca.2017a0031>
- Carey C, Gravuer K, Gennet S, et al. 2020. Supporting evidence varies for rangeland management practices that seek to improve soil properties and forage production in California. *Calif Agr* 74(2):101–11. <https://doi.org/10.3733/ca.2020a0015>
- Carey C, Weverka J, DiGaudio R, et al. 2020. Exploring variability in rangeland soil organic carbon stocks across California (USA) using a voluntary monitoring network. *Geoderma Regional* 22:e00304. <https://doi.org/10.1016/j.geodrs.2020.e00304>
- [CDFA] California Department of Food and Agriculture. 2016. Healthy Soil Program Incentives Program. www.cdfa.ca.gov/oeff/healthysoils/IncentivesProgram.html (accessed Mar. 1, 2019).
- CDFa. 2021. Healthy Soil Program Incentives Program. HSP Incentives Program Program-Level Data of Executed and Funded Grant Projects. www.cdfa.ca.gov/oeff/healthysoils/docs/HSP_Incentives_program_level_data_funded_projects.pdf (accessed Dec. 29, 2021).
- Chabbi A, Kogel-Knabner I, Rumpel C. 2009. Stabilized carbon in subsoil horizon is located in spatially distinct parts of the soil profile. *Soil Biol Biochem* 41:256–61. <https://doi.org/10.1016/j.soilbio.2008.10.033>
- Conant RT, Paustain K, Elliott E. 2001. Grassland management and conversion into grassland: effects on soil carbon. *Ecol Appl* 11:343–55. [https://doi.org/10.1890/1051-0761\(2001\)011\[0343:GMACIG\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[0343:GMACIG]2.0.CO;2)
- Dahlgren RA, Singer MJ, Huang X. 1997. Oak tree and grazing impacts on soil properties and nutrients in California oak woodland. *Biogeochemistry* 39:45–64. <https://doi.org/10.1023/A:1005812621312>
- Dass P, Houlton BZ, Wang Y, Warland D. 2018. Grasslands may be more reliable carbon sinks than forests in California. *Environ Res Lett* 13:074027. <https://doi.org/10.1088/1748-9326/13/7/074027>
- Dibblee TW. 2008. Geologic map of the Cummings Mountain and Tehachapi quadrangles. Dibblee Geology Center Map #DF-397. Santa Barbara Museum of Natural History.
- Eastburn DJ, O'Geen AT, Tate KW, Roche LM. 2017. Multiple ecosystem services in a working landscape. *PLoS ONE* 12:e0166565. <https://doi.org/10.1371/journal.pone.0166595>
- [FRAP] CAL FIRE Fire and Resource Assessment Program. 2018. California's Forests and Rangelands 2017 Assessment. <https://frap.fire.ca.gov/assessment/> (accessed Mar. 1, 2019).
- Gregory AS, Dungait JAJ, Watts CW, et al. 2016. Long-term management changes topsoil and subsoil organic carbon and nitrogen dynamics in a temperate agricultural system. *Eur J Soil Sci* 67(4):421–30. <https://doi.org/10.1111/ejss.12359>
- Herrero M, Henderson B, Havlik P, et al. 2016. Greenhouse gas mitigation potentials in the livestock sector. *Nat Clim Change* 6:452–61. <https://doi.org/10.1038/nclimate2925>
- Herrick JE, Wander MM. 1997. Relationships between soil organic carbon and soil quality in cropped and rangeland soils: the importance of distribution, composition, and soil biological activity. In *Soil Processes and the Carbon Cycle*. Lal R, Kimble JM, Follett RF, Stewart BA (eds). New York, NY: CRC Press LLC. p 405–25.
- Kaiser M, Ellerbrock RH, Wulf M, et al. 2012. The influence of mineral characteristics on organic matter content, composition, and stability of topsoils under long-term arable and forest land use. *J Geophys Res* 117:G02018. <https://doi.org/10.1029/2011JG001712>
- Koteen LE, Baldocchi DD, Harte J. 2011. Invasion of non-native grasses causes a drop in soil carbon storage in California grasslands. *Environ Res Lett* 6:044001. <https://doi.org/10.1088/1748-9326/6/4/044001>
- Latta FF. 1976. *Saga of Rancho El Tejon*. Exeter, Calif.: Bear State Books. 293 p.
- Matzek V, Stella J, Ropion P. 2018. Development of a carbon calculator tool for riparian forest restoration. *Appl Veg Sci* 21:584–94. <https://doi.org/10.1111/avsc.12400>
- Ogle SM, Breidt FJ, Easter M, et al. 2010. Scale and uncertainty in modeled soil organic carbon stock changes for US croplands using a process-based model. *Glob Change Biol* 16(2):810–22. <https://doi.org/10.1111/j.1365-2486.2009.01951.x>
- Rau BM, Johnson DW, Blank RR, Chambers JC. 2009. Soil carbon and nitrogen in a Great Basin pinyon-juniper woodland: influence of vegetation, burning, and time. *J Arid Environ* 73(4-5):472–9. <http://doi.org/10.1016/j.jaridenv.2008.12.013>
- Robertson GP, Sollins P, Ellis BG, Lajtha K. 1999. Exchangeable ions, pH, and cation exchange capacity. In *Standard Soil Methods for Long-Term Ecological Research*. Robertson GP, Coleman DC, Blendsoe CS, Sollins P (eds). New York, New York: Oxford Press. p 106–14.
- Ryals R, Kaiser M, Torn MS, et al. 2014. Impacts of organic matter amendments on carbon and nitrogen dynamics in grassland soils. *Soil Biol Biochem* 68:52–61. <https://doi.org/10.1016/j.soilbio.2013.09.011>
- Sanderson JS, Beutler C, Brown JR, et al. 2020. Cattle, conservation, and carbon in the western Great Plains. *J Soil Water Conserv* 75:5A–12A. <https://doi.org/10.2489/jswc.75.1.5A>
- Schade JD, Sponseller R, Collins SL, Stiles A. 2003. The influence of Prosopis canopies on understory vegetation: Effects of landscape position. *J Veg Sci* 14(5):743–50. <https://doi.org/10.1111/j.1654-1103.2003.tb02206.x>
- Schade JD, Hobbie S. 2005. Spatial and temporal variation in islands of fertility in the Sonoran Desert. *Biogeochemistry* 73:541–53. <https://doi.org/10.1007/s10533-004-1718-1>
- Schlesinger WH, Raikes JA, Hartley AE, Cross AF. 1996. On the spatial pattern of soil nutrients in desert ecosystems. *Ecology* 77(2):364–74. www.resolutionmineeis.us/documents/schlesinger-et-al-1996
- Schuman GE, Janzen HH, Herrick JE. 2002. Soil carbon dynamics and potential carbon sequestration by rangelands. *Environ Pollut* 116:391–6. [http://doi.org/10.1016/S0269-7491\(01\)00215-9](http://doi.org/10.1016/S0269-7491(01)00215-9)
- Silver WL, Ryals R, Eviner V. 2010. Soil carbon pools in California's annual grassland ecosystems. *Rangeland Ecol Manag* 63:128–36. <http://doi.org/10.2111/REM-D-09-00106.1>
- Spencer S, Ogle SM, Breidt FJ, et al. 2012. Designing a national soil carbon monitoring network to support climate change policy: a case example for US agricultural lands. *Greenhouse Gas Measurement Manage* 1(3-4):167–78. <https://doi.org/10.1080/20430779.2011.637696>
- Spiegel S, Bartolome JW, White M. 2016. Applying ecological site concepts to adaptive conservation management on an iconic Californian landscape. *Rangelands* 38(6):365–70. <https://doi.org/10.1016/j.rala.2016.10.002>
- Stanton CY, Mach KJ, Turner PA, et al. 2018. Managing cropland and rangeland for climate mitigation: an expert elicitation on soil carbon in California. *Climatic Change* 147:633–46. <https://doi.org/10.1007/s10584-018-2142-1>
- Swanston CW, Torn MS, Hanson PJ, et al. 2005. Initial characterization of processes of soil carbon stabilization using forest stand-level radiocarbon enrichment. *Geoderma* 128(1-2):52–62. www.osti.gov/servlets/purl/15016762
- Syswerda SP, Corbin AT, Mokma DL, et al. 2011. Agricultural management and soil carbon storage in surface vs. deep layers. *Soil Sci Soc Am J* 75(1):92–101. <https://doi.org/10.2136/sssaj2009.0414>
- Twisselman EC. 1967. A flora of Kern County, California. *Wasmann J Biol* 25:1–395.
- [USDA] US Department of Agriculture. 2018. Ecological Site Descriptions. www.nrcs.usda.gov/wps/portal/nrcs/main/national/technical/ecoscience/desc/ (accessed March 1, 2019).