

UC Agriculture & Natural Resources

California Agriculture

Title

Olive cultivars field-tested in super-high-density system in southern Italy

Permalink

<https://escholarship.org/uc/item/7038n8dv>

Journal

California Agriculture, 65(1)

ISSN

0008-0845

Author

Godini, Angelo

Publication Date

2011

Copyright Information

Copyright 2011 by the author(s). All rights reserved unless otherwise indicated. Contact the author(s) for any necessary permissions. Learn more at <https://escholarship.org/terms>

Peer reviewed

Olive cultivars field-tested in super-high-density system in southern Italy

by Angelo Godini, Gaetano Alessandro Vivaldi, and Salvatore Camposeo

According to the International Olive Oil Council, world olive oil consumption has risen from 2.8 million tons (1991-1992) to 3.5 million tons (2005-2006), due to increases in the consumption of healthier foods in many countries, including the United States. The market outlook for extra-virgin olive oil is very good, and many countries are actively increasing their olive acreages, particularly in North Africa, the Middle East, South America, Australia and the United States (Godini 2010).

The Mediterranean's traditional olive industry is based on production systems that are hundreds of years old and characterized by low yields and high production costs. The European Union subsidy system, which has helped European olive farmers to stay in busi-

ness, will end in 2014. Moreover, the application of a "free exchange" area in 2010 will legalize the importation of lower-cost extra-virgin olive oils from the southern Mediterranean Basin into Europe (Godini 2010). Year after year, the profitability of Italy's traditional olive culture becomes increasingly doubtful, notwithstanding the world-wide renown of so-called "Made in Italy" extra-virgin olive oil (Godini and Bellomo 2002).

California production of extra-virgin olive oil is reportedly about 2% of total U.S. consumption, with the rest imported mainly from Italy and Spain. In recent years, California has started increasing its oil olive acreage. California olive growers have planted more than 22,000 acres since 1999, about 12,000 acres of which is in the super-high-density olive system, with tree densities of 676 per acre or more. This system allows for mechanical planting and harvesting of olives, reducing labor costs.

We believe that super-high-density olive culture can help to assure profitability for both European and U.S. olive growers in the coming decades. This model, born in Spain at the end of the 20th century, has resulted in noticeable increases in yield per acre. Up until now, super-high-density olive culture has utilized a limited number of cultivars, primarily 'Arbequina', 'Arbosana' and 'Koroneiki', which possess suitable features such as a semi-dwarf habit, early bearing (first production at the second-to-third year after planting), consistent initial crops (more than 2.2 pounds per plant), crop stabilization between 5 and 6 years, and fruit that is impact-resistant and has good oil quality (Godini and Bellomo 2002).

The results that we present here are preliminary. Considering that Italy's Mediterranean climate is similar to California's, we believe that soil and

climate differences should have little influence on the applicability of these findings to California.

Experimental orchard

In summer 2006, we established a new experimental orchard at Valenzano, near Bari, in the experimental farm of the Dipartimento di Scienze delle Produzioni Vegetali at University Aldo Moro of Bari, Italy. In addition to standard clones of 'Arbequina',

'Arbosana' and 'Koroneiki', two additional cultivars were introduced: 'Coratina', the most popular Apulian olive oil cultivar, and 'Urano', a new Italian cultivar considered by our research group to be well-suited for super-high-density olive culture.

The olive trees were propagated in commercial nurseries by softwood cutting, and the experimental orchard was established according to the super-high-density planting scheme (676 plant per acre, with a tree spacing of 157.5 inches by 59 inches) and a north-south row orientation. The trees were trained to central leaders. Drip irrigation was supplied to each tree every 3 days between late spring and late summer, increasing from 423 cubic yards per acre annually in 2006, to 476 in 2007, to 794 in 2008 and 2009. Harvesting was performed on Nov. 20 in 2008 and 2009, in the third and fourth years after planting, respectively, using the Pellenc Activ' 4560 harvesting machine.

Cultivar performance

Vegetation. In December 2009, the average tree height had reached 107 inches, 5.3 times the initial growth of the previous year, with a maximum of 7.6 times more growth for 'Arbequina' and a minimum of 2.2 times more for 'Urano' (table 1). Only the crown width of 'Coratina' exceeded 79 inches by



Super-high-density hedgerow planting systems for olives employ over-the-row harvesters (shown, in California).

ness, will end in 2014. Moreover, the application of a "free exchange" area in 2010 will legalize the importation of lower-cost extra-virgin olive oils from the southern Mediterranean Basin into Europe (Godini 2010). Year after year, the profitability of Italy's traditional olive culture becomes increasingly doubtful, notwithstanding the world-wide renown of so-called "Made in Italy" extra-virgin olive oil (Godini and Bellomo 2002).

TABLE 1. Tree height at planting (June-July 2006) and December 2009, crown width in December 2009

Cultivar	Tree height		Crown width
	Planting	December 2009	December 2009
<i>inches</i>			
Arbequina	14.1b*	107.3b	77.5b
Arbosana	13.0b	97.0c	77.7b
Coratina	16.9b	120.9a	96.5a
Koroneiki	16.0b	117.6a	78.4b
Urano	41.1a	92.7c	74.5b
Mean	20.2	107.1	80.9

* Within the same column and for a single parameter, different letters mark values significantly different at P = 0.01 (SNK test).

Paul M. Vossen

December 2009, exceeding the harvester tunnel size and requiring pruning intervention to control its transverse canopy growth.

Annual yields. The average annual yield in the third year was 7.7 pounds per tree, equivalent to 2.3 tons per acre; only 'Urano' exhibited a surprisingly high yield of 3.7 tons per acre (table 2). In the fourth year, the average crop yield was 11 pounds per tree or 3.3 tons per acre (up 40% from 2008), and it was more variable among cultivars. 'Koroneiki', 'Arbosana' and 'Arbequina' followed by 'Coratina' gave satisfactory yield (between 13.7 and 11.3 pounds per tree or 4.1 and 3.4 tons per acre). The yield for 'Urano' dropped to 4.9 pounds per tree or 1.5 tons per acre, perhaps due to heavy cropping in the previous year.

Cumulative yields and oil. We also compared cumulative yields over the first 4 years of the trial. 'Koroneiki' showed the highest cumulative yield (41.3 tons per acre), and 'Urano' was relatively less productive (31.6 tons per acre) (table 2). The peculiar behavior of 'Urano' requires further investigation. Considering its average overall oil content of about 17.0%, 'Koroneiki' was the most productive cultivar with 6.9 tons per acre of oil over 4 years. The other cultivars exhibited similar cumulative oil production.

Harvesting efficiency, fruit and shoot damage. Harvesting efficiency was satisfactory on the whole (93.1%), notwithstanding differences among cultivars (table 3). 'Arbequina', 'Coratina' and 'Urano' had the highest harvest

TABLE 2. Fruit production per year, cumulative yield at the third (2008) and fourth (2009) year after planting, and mean oil output and cumulative production

Cultivar	Fruit production		Cumulative yield*	Mean oil output	Cumulative oil production
	3rd year	4th year			
 pounds/tree† tons/acre		tons/acre	%	tons/acre
Arbequina	5.5b‡	12.4ab	33.4b	17.7	5.68b
Arbosana	6.2b	12.6ab	35.1ab	17.5	5.83b
Coratina	7.1b	11.3b	34.1b	17.3	5.86b
Koroneiki	8.4b	13.7a	41.3a	15.0	6.89a
Urano	12.1a	4.9c	31.6c	16.8	5.26b
Mean	7.7	11.0	35.1	16.9	5.91

* Over four years of the trial.

† 7.73 pounds per tree = 2.3 tons per acre.

‡ Within the same column and for a single parameter, different letters mark values significantly different at $P = 0.01$ (SNK test).

efficiency; 'Arbosana' and 'Koroneiki' were less satisfactory. But these differences were due to the fruit-ripening stages reached by each cultivar at the harvesting date: mature for 'Arbequina', 'Coratina' and 'Urano', but immature 'Arbosana' and 'Koroneiki'.

No damaged fruits were reported for 'Arbequina', 'Arbosana' and 'Koroneiki', whereas 'Coratina' and 'Urano' exhibited very low percentages of damaged fruit.

The average percentage of shoots damaged per tree by the harvesting machine beaters was insignificant at less than 1.0%. Of these damaged shoots, young and thin current-year shoots incurred the most damage (80.0%), perhaps because they were more exposed. The percentage of damaged shoots up to 1 inch in diameter was 14.3%, and to shoots thicker than 1 inch was 5.7%. Only 'Coratina' and 'Urano' exhibited a significant percentage of broken shoots or branches thicker than 1 inch: 'Coratina' because of its spreading habit between rows, and 'Urano' because of its spreading habit and thick, bending branches.

High density, high yields

The present data confirms and improves upon results obtained in previous experimental trials (Camposeo and Godini 2010). In terms of early bearing and yield consistency, all the tested cultivars performed satisfactorily. And in sensory evaluations, the resulting extra-virgin oils had sweet typology and were well-balanced, highly fruity and ready to use (Camposeo et al. 2010).

We know that higher yields have been recorded elsewhere with super-high-density olive culture; however, we consider annual yields of about 17.5 tons per acre of fruit to be satisfactory. In fact, this value, equivalent to a yield of only about 9.4 pounds per tree, would be helpful in avoiding alternate bearing and subsequent problems that could cause conflicts between vegetative growth and cropping consistency. Tree size can be controlled by pruning when they grow larger than the size of the harvester head. Our data indicates that the noted yield limit was reached by at least four out of five cultivars in just the 4th year after planting.

A. Godini is Professor, G. Allesandro Vivaldi is Ph.D. Student, and S. Camposeo is Ph.D. Researcher, Dipartimento di Scienze delle Produzioni Vegetali, University Aldo Moro of Bari, Italy. The authors wish to thank Provincia di Bari, Italy, for its financial support of this research.

References

- Camposeo S, Ferrara G, Palasciano M, Godini A. 2008. Varietal behaviour according to the superintensive olive-culture training system. *Acta Hort* 791:271-4.
- Camposeo S, Godini A. 2010. Preliminary results about the performance of 13 cultivars according to the super high density olive-culture training system in Apulia (Southern Italy). *Adv Hort* 24(1):16-20.
- Camposeo S, Vivaldi GA, Gallotta A, et al. 2010. Valutazione chimica e sensoriale degli oli di alcune varietà di olivo allevate in Puglia con il modello superintensivo. *Frutticoltura* 72(6):80-83 (in Italian with summary in English).
- Godini A. 2010. L'olivicultura italiana tra valorizzazione e innovazione. *Frutticoltura* 72(6):52-69 (in Italian with summary in English).
- Godini A, Bellomo F. 2002. Olivicultura superintensiva in Puglia per la raccolta meccanica con vendemmiatrice. International Congress of Oliveculture. Spoleto (Italy), April 22-23:230-4 (in Italian with summary in English).

TABLE 3. Harvesting efficiency, damaged fruits and damaged shoots per tree, as mean of the third and fourth year after planting

Cultivar	Harvesting efficiency	Fruit damaged	Shoots damaged
 %		n/tree
Arbequina	97.2a*	0.0c	0.4b
Arbosana	87.1b	0.0c	0.5ab
Coratina	97.3a	1.5a	0.2c
Koroneiki	87.6b	0.0c	0.6a
Urano	96.0a	0.4b	0.2c
Mean	93.1	0.4	0.4

*Within the same column and for a single parameter, different letters mark values significantly different at $P = 0.01$ (SNK test).