

UC Merced

Biogeographia - The Journal of Integrative Biogeography

Title

Faunistics and zoogeographical overview of the Mediterranean and Black Sea marine Gastrotricha

Permalink

<https://escholarship.org/uc/item/95b3x62x>

Journal

Biogeographia - The Journal of Integrative Biogeography, 24(1)

ISSN

1594-7629

Authors

Todaro, M. Antonio

Matinato, Lucia

Balsamo, Maria

et al.

Publication Date

2003

DOI

10.21426/B6110169

Peer reviewed

Faunistics and zoogeographical overview of the Mediterranean and Black Sea marine Gastrotricha

M. ANTONIO TODARO*, LUCIA MATINATO**,
MARIA BALSAMO***, PAOLO TONGIORGI**

*Dipartimento di Biologia Animale, Università di Modena e Reggio Emilia,
Via Campi 213/d, I-41100 Modena (Italy);
e-mail: todaro.antonio@unimo.it

**Dipartimento di Scienze Agrarie, Università di Modena e Reggio Emilia,
Via Kennedy 17, I-42100 Reggio Emilia (Italy)

***Istituto di Scienze Morfologiche, Università di Urbino,
Via Oddi 21, I-61029 Urbino (Italy)

Key words: Gastrotrichs, Biogeography, Meiofauna, Taxonomy, Mediterranean Sea, Black Sea.

SUMMARY

The paper provides an updated overview of the knowledge regarding the marine gastrotrich fauna of 10 Mediterranean and 2 Black Sea countries. Taxonomic account and species distribution come from published records, including electronically disseminated information, as well as original data from 18 localities (16 western and 2 levantine), investigated by the authors between 1990 and 2002. Since the description of the first marine Mediterranean gastrotrich, *Hemydys agaso* Claparède, 1867, discovered in the Gulf of Naples (Italy), 273 additional species, from 417 locations have been recorded in the basin so far. Of the gastrotrichs found, 143 species, in 24 genera and 6 families, belong to the order Macrodasyida and 131 species, in 11 genera and 3 families, belong to the order Chaetonotida; these statistics include several species as yet not described. The number of species per location is variable, ranging from 1-37, with a global mean of 8.04 ± 6.69 spp/location. Egypt and Israel show mean values well above the average, with 12.8 and 10.8 spp/loc. respectively, whereas mean values for Algeria, Tunisia, Bulgaria, Croatia, and France are below average (1-5.43 spp /location); data for Cyprus, Greece, Romania and Italy are within the average value. *Acanthodasys aculeatus*, recorded in 28% of the investigated locations is the most frequently found macrodasyidan; *Halichaetonotus acutifer*, also recorded in 28% of the investigated localities, is the most common chaetonotidan. Data analysis indicated substantial differences among countries, regarding sampling effort and, consequently, faunistic knowledge. In comparison with the generally good information concerning the Italian fauna (177 species from 246 localities), gastrotrichs from other Mediterranean nations are much less known; along with Italy, only few other Nations have been investigated to a sufficient extent (i.e., Greece: 44 sampled localities, 77 recorded species; France: 37 L, 70 spp; Egypt: 28 L, 81 spp; Israel: 15 L, 55 spp; Cyprus: 9 L, 41 spp), whereas investigation carried out in several other countries can be considered, to a variable degree, incomplete (i.e., Algeria: 16 investigated localities and 11 species found; Romania: 7 L, 30 spp; Tunisia: 5 L, 2 spp; Bulgaria: 3 L, 11 spp; Croatia: 1 L, 12 spp; Spain: 1 L, 3 sp, Malta: 2L, 2 spp.), or nil (e.g., Morocco, Libya, Turkey, Albania). All marine gastrotrich families and most marine genera have representatives in the Mediterranean fauna, notable absences regard the genera *Desmodays*, *Dinodays*, *Planodays*, *Prostobuccania* and perhaps *Pseudoturbanella*; on the other hand representatives of two genera, *Emydays* and *Dendropodola*, are unknown elsewhere in the world except the western Mediterranean. At species level, about 140 taxa have so far been found only in the Mediterranean area, whereas the

remainder have some north-Atlantic connections, including several considered as regional cosmopolitans (i.e., reported from at least two Oceans). Within the Mediterranean, the species geographic distribution does not appear to be homogeneous, with several taxa restricted to either the western or to the Levantine basin; yet, Italy, Egypt and Israel show the highest numbers of endemic species, 70, 24 and 14 respectively, whereas only one species seems restricted to the Black Sea. A cluster analysis using the Bray-Curtis presence-absence similarity value to examine the relationship between 11 geographic areas, shows a clear separation between the Black Sea and the Mediterranean regions and within this, the Algerian-Tunisian region is separated from and the remaining 9 areas. The latter appear arranged in two main clusters, one including the southern Levantine regions (i.e., Egypt, Israel and Cyprus), the other containing regions of the north-western basin, the Adriatic Sea and the two Greek sub-regions. In this cluster, the region comprising coastal France and Formentera (Spain) appears separated from the one containing, arranged into two distinct subsets, the Italian and the Greek areas. However, due to the paucity of information, data from key-areas is urged to confirm or disprove our current perception about numbers and distribution of gastrotrich taxa within the Mediterranean and the Black Sea.

INTRODUCTION

Gastrotricha are a phylum of aquatic, free-living metazoans adding up to about 690 microscopic species, ranging from 80 µm to 3 mm in total length. Taxonomically, gastrotrich species are split into two orders, the marine Macrodasyida (6 families, 23 genera), and the marine and freshwater Chaetonotida (7 families, 28 genera) showing great differences from morphological to reproductive points of view. Macrodasyidans are characterised by a worm-like shape and the presence of numerous anterior, lateral and posterior (caudal) adhesive tubes; they are hermaphrodites, mainly simultaneous or sequential. In contrast, chaetonotidans are smaller and characterised by a tenpin body shape having a furcate posterior end, called the furca, which terminates with the only two adhesive tubes exhibited by these animals (except *Neodasys*). Chaetonotidans are mainly parthenogenetic, with hermaphroditic species restricted to selected taxa (e.g., *Musellifer*, *Neodasys*, Xenotrichulidae).

In marine settings, Gastrotricha represent an ever present component of the meiobenthic community of the sandy substrata, often ranking third in abundance following Nematoda and Harpacticoida Copepoda (e.g., Coull, 1985). They colonize a variety of habitats, from fine to coarse sand in both littoral and sublittoral areas and also caves; however, abundance and biodiversity peak in medium, clean sand of the shallow subtidal areas (Todaro and Rocha, 2003). Representative of both orders move between sand grains by means of ventral locomotory cilia, occasionally attaching to the substratum by means of adhesive tubes; like other meiobenthic animals, gastrotrichs have no planktonic dispersal stages so juveniles undergo direct development in the same habitat as their parents. Based of bio-physiological traits such as short life span, small number of offspring, absence of dispersal stages, limited swimming ability of the adults, gastrotrich species would be expected to have restricted geographic range; however, in contrast with this prospect, many species are reported having a wide geographic range, with several being amphi-atlantic or even cosmopolitan. Research addressing this apparent paradox has undergone a recent revival due to both theoretical and technical advances (Todaro et al., 1996; Todaro and Rocha, 2003). In this framework, the aim of the present study is to provide a

handy, yet exhaustive overview, of the gastrotrich fauna of one of the best known basins in the world, as contribution to debate on the biogeography of these acoelomate worms. The complimentary pictorial keys of the gastrotrich genera found in the Mediterranean basin will hopefully prove useful not only to gastrotrich specialists but also to marine ecologists who find these abundant metazoans in the course of research on interstitial meiobenthos.

MATERIAL AND METHODS

Taxonomic account and species distribution come from published records, including electronically disseminated information (cf. Hummon 2001a), as well as original data from 18 new localities (16 in the western and 2 in the eastern basin), investigated by the authors between 1990 and 2002 time frame. Below is a list of the consulted papers, arranged by nation, containing taxonomic and/or faunistic data; in brackets is the number of papers per nation:

ALGERIA (1): d'Hondt (1974); TUNISIA (1): Westheide (1972); MALTA (1): Boaden (1965a); BULGARIA (1): Valkanov (1957); ROMANIA (3): Rodewald (1938); Rudescu (1966, 1967); GREECE (1) Hummon and Roidou (1994); CROATIA (1): Schrom (1972); ITALY (51): Balsamo and Todaro (1987); Balsamo et al. (1992, 1994, 1995, 1996, 1997); Boaden (1965a,b); Claparède (1867); Clausen (1965); De Zio and Grimaldi (1964); De Zio (1965); Evans et al. (1993); Faienza et al. (1999, 2000); Fregni (1998a); Fregni et al. (1998, 1999); Gerlach (1953); Grünspan (1908); Hummon et al. (1990a, 1992, 1993, 1996, 1998); Luporini et al. (1971, 1972; 1973a,b); Papi (1957); Remane (1927a,b, 1951, 1952); Renaud-Mornant (1968); Schrom (1966a,b, 1972); Todaro (1992, 1997, 1998a,b); Todaro and Balsamo (1994); Todaro et al. (1988, 1992, 2001, 2003); Tongiorgi (1975); Tongiorgi and Balsamo (1984); Tongiorgi et al. (1999); Wilke (1954); FRANCE, including Corsica: (9) Delamare-Deboutteville (1953, 1954); Fize (1957, 1963); Hummon et al., (1992); Remane (1961); Swedmark (1956); Todaro et al. (2002); Vivier (1974).

Information about Egypt, Israel and Cyprus, are from Hummon's (2001a) CD-rom "Global database for marine Gastrotricha", which is available from the author: "hummon@ohio.edu" or in second place from us: "todaro.antonio@unimo.it"; Hummon's CD-rom also supplied additional information from Greece and Mediterranean France and was also used to cross-check data gathered from literature. For general information about the CD-rom and the source of data it contains see Hummon (2001b,c) and the section "Results and Discussion" below.

Original records come from 18 locations (13 Italian, 2 Greek, 1 French, 1 Spanish, Table I); samples were collected and processed according to our standardised methods (e.g., Todaro et al., 2002). Global data (old + new records) were arranged in a species by nation *check-list* (see appendix I and II).

Since Hummon (2001a) also includes in his CD-rom a conspicuous number of putative new species, as yet undescribed, we have incorporated these taxa in the list. We trust, in doing so, to hopefully give a more realistic (better) picture about the biodiversity and zoogeography of Mediterranean gastrotrich-fauna. Likewise, most of the unnamed species, putatively new to science, reported in Todaro et al. (2001a) appear also in the present check-list, with the exception of those for which information granting them a clear distinctness, is not considered sufficient anymore.

The Cluster analysis performed on a species by location matrix was carried out using the software Primer 5.2.0. Prior analysis, data were pooled in the following 11 main areas: 1, Algeria + Tunisia; 2, Egypt; 3, Israel; 4, Cyprus; 5, Black Sea (= Romania + Bulgaria); 6, Greece-Aegean (= Attica + Cyclades + Macedonia + Peloponnesis + Sporades + Thessaly); 7, Greece-Mediterranean (= Crete + Dodecannese); 8, It-Adriatic Sea (= Adriatic coast of Italy, from Trieste to Santa Maria di Leuca + Rovjni in Croatia); 9, It-Ionian Sea (= Italian Ionian coast of Apulia, Lucania, Calabria and Sicily); 10, It-Tyrrhenian Sea (= western coast of Italy, including Sardinia and the northern coast of Sicily + Corsica); 11, France-Spain (Continental coast of France + a single site on the island of Formentera, Spain).

RESULTS AND DISCUSSION

Literature data

First record of a marine gastrotrich species from the Mediterranea Sea dates back to the XIX century and comes from Claparède (1867) who, investigating the minute benthic organisms of the Gulf of Naples, finds and describes *Hemidasys agaso*, a species affiliated later on with the Thaumastodermatidae. During the first half of the XX century only three other papers dealing with Mediterranean gastrotrichs were published: Grünspan (1908) describes two species of *Ichthydium* from the Gulf of Trieste, whereas Remane (1927a,b) reports five new species (*Dactylopodola typhle*, *Lepidodasys platyurus*, *Macrodasys caudatus*, *Tetranchyroderma megastomum* and *Thaumastoderma mediterraneus*), and four new genera and species (*Paraturbanela dohrni*, *Platydasys maximum*, *Ptychostomella mediterranea*, *Diplodasys platydasyooides*) from the Gulf of Naples. At about the same time Rodevald (1938) reports the first species from the Romanian coast of the Black Sea. A formidable impulse to the study of the Mediterranean gastrotrich fauna was given by German researchers during 1950s: Remane (1951, 1952) describing two new species from Cuma (Italy), and Sete (France) respectively, Gerlach (1953) who worked along the coast of




Fig. 1 - Representative species belonging to families and genera found in the Mediterranean basin. A-L: Chaetonotidae, A: *Aspidiophorus polystictos*; B: *Chaetonotus apechochaetus*; C: *C. dispar*; D: *C. lacunosus*; E: *Halichaetonotus aculifer*; F: *Halichaetonotus margaretae*; G: *Heterolepidoderma istrianum*; H: *Ichthydium tergestinum*; I: *Lepidodermella squamata*; L: *Musellifer delamarei*.

Tuscany, and especially Wilke (1954) whose doctoral work on the fauna from the Gulf of Naples doubled the number of species known for the Mediterranean, which raised to 43. During the same decade important contributions were made by Swedmark (1956) and Fize (1957) investigating the gastrotrich fauna of the southern coast of France, by Papi (1957) who described three new species from the surroundings of Naples, and by Valkanov (1957) who, working along the Bulgarian coast, published the second out of the only three papers on gastrotrichs of the Black Sea, the third being that of Rudescu (1966) devoted again to the gastrotrichs of Romania. Between the 60s and 70s of the past century, key information came from the taxonomic work of Fize (1963) in France, Boaden (1965a,b) in Italy and Malta, Clausen (1965), Schrom (1966a,b) and Renaud-Mornant (1968) in Italy, Schrom (1968) in Croatia, and by the ecological work of de Zio and Grimaldi (1964) and de Zio (1965) carried out at some Apulian beaches of Italy. A flourishing of papers characterised the two decades between 1970-1990, some of the more extensive work focused on regional faunas, e.g., Italy: coast of Tuscany (Luporini et al., 1971, 1973a,b; Tongiorgi, 1975), northern Adriatic Sea (Schrom, 1972); Algeria: (d'Hondt, 1974); others dealt with the description of single species (Vivier, 1974; Tongiorgi and Balsamo, 1984; Balsamo and Todaro 1987, Todaro et al., 1988). During the same time-frame records of nominal species also appeared in papers not specifically focused on Gastrotricha, this data is important mostly because it was gathered from areas never sampled again (e.g., Tunisia, Westheide, 1972). Consequently, by the end of the 80s of the last century the gastrotrich fauna of the Mediterranean Sea was already one of the best known world wide (with more than 90 known species). New and more intensive research was carried out between 1990 and 2002 headed by a team of American-Italian investigators whose goal was two fold: a) to verify previous data (mainly resampling locations already investigated), and b) to add information from poor known areas. The standardization of the methodology, implemented by new and better equipment (i.e., light microscopes equipped with DIC optics, scanning electron microscopy, videomicroscopy, videorecording etc.) would have made possible the building-up of a coherent data-set, useful for several purposes, from specimen comparisons to biogeographic analysis. Research performed along the Italian coasts spawned a number of papers which focused on taxonomy and ecology (Hummon et al., 1990a, 1992, 1993, 1996, 1998; Balsamo et al., 1992, 1994, 1995, 1996, 1997; Todaro, 1992, 1997, 1998a,b; Evans et al., 1993; Todaro et al., 1992; Todaro and Balsamo, 1994; Fregni, 1998a; Fregni et al., 1998, 1999; Faienza et al., 1999, 2000; Tongiorgi et al., 1999), and provided live material to perform studies aimed at shedding light on general topics such as reproductive biology (e.g., Balsamo and Todaro, 1987; Ferraguti and Balsamo, 1994, 1995; Ferraguti




Fig. 2 - Representative species belonging to families and genera found in the Mediterranean basin. A-C: Xenotrichulidae, A: *Draculiciteria tesselata*; B: *Heteroxenotrichula squamosa*; C: *Xenotrichula intermedia*. D: Neodasyidae, D: *Neodays uchidai*. E-H: Dactylopodolidae, E: *Dactylopodola mesophyle*; F: *Dendrodasys affinis*; G: *Dendropodola transitionalis*; H: *Xenodasy sancti-gouveni*, a species from the Atlantic coast of France (Roscoff), morphologically close to *Xenodasy* sp. found along the Ionian coast of Italy. I-N: Lepidodasyidae, I: *Cephalodasy turbanelloides*; L: *Dolichodasy elongatus*; M: *Lepidodasy martini*; N: *Megadasy minor*.

et al., 1995; Fregni, 1998b; Balsamo et al., 2002) life history (Balsamo and Todaro, 1988), and even on phylogeny (Todaro et al., 1996; Wirz et al., 1999; Todaro et al., 2003). A checklist of the Italian gastrotrich-fauna including both marine and freshwater forms was published by Balsamo and Tongiorgi (1995) and has been updated in the *Inventory of Italian marine gastrotrichs: an annotated check list* by Todaro et al. (2001a) for the marine species.

Written accounts of studies carried out outside of Italy, dealing with marine gastrotricha from Greece and Corsica are reported in Hummon and Roidu (1994) and Todaro et al. (2002) respectively. In the outside-Italy scenarios a particular relevance bear the extensive research carried out by W. D. Hummon in the Levantine countries (i.e., Greece, Israel, Egypt, Cyprus) in collaboration with several local researchers, and in the Roussignol coast of France (together with M. A. Todaro) between 1990-2000. Information on this research has been published in several abstracts (Hummon et al., 1990b; Hummon and Hummon, 1992, 1995a,b; Hummon et al., 1994) and reported in a more extended layout in the *Global database for marine Gastrotricha* (Hummon, 2001a). It is a useful tool that the author generously makes available, on request, to the scientific community (see Materials and Methods above). In a larger framework the “The Global database” by Hummon (2001a) furnishes an ample and authoritative source of information regarding the world geographical distribution of all marine gastrotrichs species known until 2001.

New data

Of the 18 new investigated sites, 14 are along the Italian coasts, two are located in Crete (Greece), and one in each of Spain (Formentera, Baleari Islands) and France. Geographical and ecological information about localities are given in Tables I and II whereas reported in Tables III and IV are the taxonomic account and the distribution of the recorded species. Of the 48 species found, *Tetranchyroderma inaequitubulatum* is new to the Italian fauna and its record is the second ever for the species; *Mesodasys adenotubulatus*, *T. esarabdophorum* and *T. thysanophorum* are new to Greece; *Mesodasys adenotubulatus*, *Paraturbanella teissieri* and *Heteroxenotrichula squamosa* represent the first records of marine gastrotrich species from Mediterranean Spain.

Global data

All together 274 species (3046 records, i.e., species × sites) have been reported from 416 locations in 13 countries. Of these, 143 species, in 24 genera and 6 families, belong to the order Macro dasyida (Appendix I, Figs. 2-4), and 131 species, in 11 genera and 3 families, belong to the order




Fig. 3 - Representative species belonging to families and genera found in the Mediterranean basin. A-C: Lepidodasyidae, A: *Mesodasys adenotubulatum*; B: *Paradasys subterraneus*; C: *Pleurodasys helgolandicus*. D,E: Macrodasyidae, D: *Macrodasys thucus*; E: *Urodasys viviparus*. G: Planodasyidae, F: *Crasiella diplura* a species from the North Sea, morphologically close to *Crasiella* sp. found along the Italian shores. G,H: Thaumastodermatidae, G: *Acanthodasys aculeatus*; H: *Diplodasys ankeli*.

Tab. I - Eighteen new, or newly investigated Mediterranean locations; geographic coordinates, date of collection, water depth, salinity and temperature of the water at the sublittoral sites.

Location	Lat. (North)	Long. (East)	Date	Salinity (‰)	Temp. (°C)	Depth (m)
Italy						
1. Cala Nave (Island of Ventotene)	40° 47'	13° 26'	29/05/01	37.5	22	3.5
2. Porto S. Felice a Circeo (Latium)	41° 13'	13° 05'	29/05/01	36.5	22	2.0
3. Fiumicino* (Latium)	41° 47'	12° 13'	12/12/03	37	15	Na
4. Marina di Alberese	42° 39'	11° 04'	30/05/01	38	22	2.0
5. Pomonte (Elba Island)	42° 47'	10° 06'	3/11/01	38	20	12.0
6. San Teodoro (Sardinia)	40° 47'	09° 41'	20/06/01	35	21	1.5
			11/06/02	36	20	2.5
7. Baia di Anzachena (Sardinia)	41° 08'	09° 28'	11/06/02	36	21	3.0
8. Portu la Rena, Castelsardo (Sardinia)	40° 55'	08° 43'	08/06/02	38	20	13.0
9. Palau (Sardinia)	41° 12'	09° 20'	09/06/02	37	21	3.0
10. Spiaggia della Mormorata (Sardinia)	41° 13'	09° 17'	07/06/02	36	20	2.0-3.0
11. San Isidoro (Apulia)	40° 09'	17° 57'	03/07/02	36	25	3.0
12. Baia Verde (Apulia)	40° 00'	18° 01'	03/07/02	38	25.7	-
13. Brindisi (Apulia)	40° 31'	18° 05'	21/09/02	37	24	2.0
14. Ravenna Lido (Emilia Romagna)	44° 28'	12° 18'	20/5/01	30	22	1.5
Greece						
15. Matala (Crete, south side)	34° 59'	24° 45'	10/09/00	38	25	2.0-4.0
16. Amnissos (Crete, north side)	35° 20'	25° 16'	11/09/00	37	26	2.0-3.0
Spain						
17. Formentera (Baleari islands)	38° 41'	01° 28'	28/07/01	35.4	24	2.0
France						
18. Cannes	43° 33'	06° 59'	10/08/01	37	25	2.0

*, data refers to littoral site; Na, not applicable.

Chaetonotida (Appendix II, Figs. 1, 2). The highest species richness is in Italy, with a total of 176 different species, followed by Egypt, 81 species; Greece, 80 species; France, 70 species and Israel, 55 species. The remainder of the countries showed a range between 2-42 species (Table V). *Acanthodasyus aculeatus*, recorded in 28% of the investigated locations is the most frequently found macrodasyidan whereas *Halichaetodonotus aculifer*, also recorded in 28% of the investigated localities, is the most common chaetonotidan (Fig. 5A,B).

The number of species per location is variable, ranging from 1-37, with a global mean of 8.04 ± 6.69 spp/location. Egypt and Israel show mean values well above the average, with 12.8 and 10.8 spp/loc. respectively, whereas mean values for Algeria, Tunisia, Bulgaria, Croatia, and France are below average (1- 5.43 spp /location); data for Cyprus, Greece, Romania and Italy are within the average value (Table VI).

Tab. II - Granulometric characteristics of the sublittoral sediment of the 18 new investigated Mediterranean locations.

Location	Mean grain size (phi), and size class	Sorting (phi), and sorting class	Skewness	Kurtosis
1.	1.95, medium sand	0.7, moderately sorted	-0.18	3.16
2.	2.03, very fine sand	0.75, very well sorted	-0.87	3.33
3.	1.01, medium sand	0.90, moderately sorted	-1.04	4.58
4.	NA	NA	NA	NA
5.	1.03, medium sand	0.97, moderately sorted	0.82	3.58
6.	2.30, fien sand	0.58, moderately well sorted	-1.67	6.38
7.	1.05, medium sand	1.02 poorly sorted	0.149	2.13
8.	1.53, medium sand	0.42, well sorted	0.29	5.53
9.	0.88, coarse sand	0.81, moderately sorted	-0.161	2.59
10.	2.23, fine sand	0.6, moderately well sorted	-0.64	3.77
11.	2.11, fine sand	0.73, moderately sorted	-1.19	4.84
12.	NA	NA	NA	NA
13.	2.46, very fine sand	0.77, moderately sorted	-0.99	5.16
14.	2.85, very fine sand	0.82, moderately sorted	1.38	4.82
15.	2.09, fine sand	0.67, moderately sorted	-0.68	3.69
16.	2.29, fine sand	0.59, moderately sorted	-0.65	4.15
17.	1.79, medium sand	0.63, moderately well sorted	-0.5	2.59
18.	NA	NA	NA	NA

Na, not available.

Data analysis indicate substantial differences among countries, regarding sampling effort and, consequently, faunistic knowledge. In comparison with the generally good information concerning the Italian fauna (177 species from 246 localities), gastrotrichs from other Mediterranean nations are much less known. Taking into account the extent of the coastline, beside Italy, only few other countries have in fact been investigated to a sufficient extent (i.e., Greece: 44 sampled localities, 77 recorded species; France: 37 L, 70 spp.; Egypt: 28 L, 81 spp.; Israel: 15 L, 55 spp.; Cyprus: 9 L, 41 spp.). Investigations carried out in the other countries are, to a variable degree, incomplete (i.e., Algeria: 16 investigated localities and 11 species recorded; Romania: 7 L, 30 spp; Tunisia: 5 L, 2 spp.; Bulgaria: 3 L, 11 spp.; Croatia: 1 L, 12 spp.; Spain: 1 L, 3 sp.; Malta: 2 L, 2 spp.), or nil (e.g., Morocco, Libya, Turkey, Albania).

All marine gastrotrich families and most marine genera have representatives in the Mediterranean fauna, notable absences regard the genera *Desmodasys*, *Dinodasys*, *Planodasys*, *Prostobuccantia* and *Pseudoturbanella*; on the other hand representatives of two genera, *Hemydasys* and *Dendropodola*, are unknown elsewhere in the world except the western Mediterranean (Figs. 1-4); it is worth noting however that both excluded and endemic genera includes only 1 or 2 (*Desmodasys*) rare species, having a limited geographic distribution.

Tab. III - Gastrotricha Macrodasyida species list and distribution in 18 new investigated Mediterranean locations. Data refers to sublittoral sites, except location # 4 Fiumicino, where only the littoral site was sampled.

Taxon	Location																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
DACTYLOPODOLIDAE																		
<i>Dactylopodola mesotiphle</i>	+	-	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-
<i>Dendrodasys affinis</i>	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-
MACRODASYIDAE																		
<i>Macrodasys caudatus</i>	+	-	-	-	+	+	-	-	+	+	-	-	-	-	-	-	-	-
<i>Urodasys acanthostylis</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Urodasys viviparous</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
LEPIDODASYIDAE																		
<i>Dolichodasys elongatus</i>	-	-	-	-	-	-	-	-	+	+	+	-	-	-	-	-	-	-
<i>Lepidodasys martini</i>	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Lepidodasys unicarenatus</i>	-	-	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-
<i>Mesodasys adenotubulatus</i>	-	-	-	-	-	-	-	-	+	-	-	-	-	+	-	+	-	-
<i>Mesodasys laticaudatus</i>	-	-	-	-	+	-	-	+	+	-	+	-	-	-	-	-	-	-
PLANODASYIDAE																		
<i>Crasiella</i> sp. a	-	-	-	-	+	-	-	-	-	+	-	-	-	-	-	-	-	-
THAUMASTODERMATIDAE																		
<i>Acanthodasys aculeatus</i>	+	-	-	-	+	+	-	-	-	+	-	-	-	+	-	-	-	-
<i>Diplodasys ankelii</i>	-	-	-	-	+	+	-	+	-	-	-	-	-	-	-	-	-	-
<i>Pseudostomella etrusca</i>	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Pseudostomella roscoffita</i>	-	-	-	-	-	-	-	-	-	+	-	-	-	+	-	-	-	-
<i>Ptychostomella tyrrhenica</i>	+	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tetranch. cirrophorum</i>	-	-	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-
<i>Tetranch. esarabdomorphum</i>	-	-	-	-	-	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Tetranchyoderma hirtum</i>	-	-	-	-	+	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Tetranch. inaequitubulatum</i>	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tetranch. megastomum</i>	-	-	-	-	-	+	-	+	+	-	-	-	-	-	+	-	-	-
<i>Tetranchyoderma papii</i>	-	-	+	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tetranc. quadridentaculatum</i>	-	-	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	-
<i>Tetranch. thysanophorum</i>	-	-	-	-	+	+	-	-	+	-	-	-	-	-	+	-	-	-
<i>Tbaumastoderma heideri</i>	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tbaumast. mediterraneus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tbaumast. ramuliferum</i>	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-
TURBANELLIDAE																		
<i>Paraturbanella dhorni</i>	-	-	-	-	+	-	-	-	-	+	+	-	-	-	-	-	-	-
<i>Paraturbanella pallida</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Paraturbanella teissieri</i>	+	-	-	-	-	+	+	-	+	-	+	-	-	-	+	+	+	+
<i>Turbanella cornuta</i>	+	-	+	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-

Species of five, elsewhere widely distributed, genera have so far been found only in the western basin: i.e., *Crasiella* (1 sp.), *Paradasys* (1 sp.), *Pleurodasys* (1 sp.) *Neodasys* (1 sp.) and *Musellifer* (2 spp.); however, giving the occasionality of these findings the presence also in the eastern basin of these genera cannot be excluded. Likewise the apparent absence from the eastern basin of the genus

Tab. IV - Gastrotricha Chaetonotida species list and distribution in 18 new investigated Mediterranean locations. Data refers to sublittoral sites, except location # 4 Fiumicino, where only the littoral site was sampled.

Taxon	Location																	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
XENOTRICHULIDAE																		
<i>Daculiciteria tesserata</i>	-	-	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Heteroxenotrichula squamosa</i>	+	-	+	-	-	-	-	-	-	-	-	-	-	+	+	+	-	-
<i>Xenotrichula intermedia</i>	-	-	+	-	-	-	-	-	-	+	-	-	+	-	-	-	-	-
<i>Xenotrichula punctata</i>	+	-	-	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-
CHAETONOTIDAE																		
<i>Aspidiop. mediterraneus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Aspidiop. paramediterraneus</i>	+	-	-	-	+	-	-	+	-	-	+	-	-	-	-	-	-	-
<i>Chaetonotus aechochaetus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chaetonotus atrax</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chaetonotus lacunosus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+
<i>Chaetonotus neptuni</i>	+	-	-	-	-	+	-	+	-	-	-	-	-	+	-	-	-	-
<i>Chaetonotus sicilensis</i>	-	-	-	-	-	+	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chaetonotus variosquamatus</i>	-	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Halichaeatonotus aculifer</i>	-	-	+	-	-	+	-	-	-	-	-	-	-	+	+	-	-	-
<i>Halichaeatonotus decipiens</i>	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Halichaeatonotus paradoxus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Halichaeatonotus spinosus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Musellifer delamarei</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Tab. V - Investigated locations and gastrotrich-fauna in 13 Mediterranean and Black Sea countries.

Country	n. Location	n. Species (C, M)	n. Families (C, M)	n. Genera (C, M)
Algeria	16	11 (4C, 7M)	6 (2C, 4M)	9 (3C, 6M)
Tunisia	5	2 (0C, 2M)	1 (0C, 1M)	1 (0C, 1M)
Malta	2	2 (0C, 2M)	2 (0C, 2M)	2 (0C, 2M)
Egypt	28	81 (47C, 34M)	7 (2C, 5M)	19 (7C, 12M)
Israel	15	55 (37C, 18M)	7 (2C, 5M)	15 (7C, 8M)
Cyprus	9	42 (25C, 17M)	7 (2C, 5M)	20 (7C, 13M)
Bulgaria	3	11 (6C, 5M)	6 (2C, 4M)	7 (4C, 3M)
Romania	7	30 (16C, 14M)	7 (2C, 5M)	16 (7C, 9M)
Greece	46	80 (45C, 35M)	7 (2C, 5M)	20 (7C, 13M)
Croatia	1	12 (9C, 3M)	4 (2C, 2M)	7 (4C, 3M)
Italy	246	177 (70C, 107M)	8 (2C, 6M)	34 (10C, 24M)
France*	37	70 (27C, 43M)	8 (3C, 5M)	24 (8C, 16M)
Spain	1	3 (1C, 2M)	3 (1C, 2M)	3 (1C, 2M)
Global data	417	274 (131C, 143M)	9 (3C, 6M)	35 (11C, 24M)

*including Corsica; C, Chaetonotida; M, Macrodayida.

Platydasy, particularly speciose in the western region (but present also in the Black Sea), seems due more to research shortfalls than to stringent eco-physiological requirement of these species, or to the geological history of the Mediterranean area.

Tab. VI - Statistics about number of species per location, most common species of the two orders found in 13 Mediterranean and Black Sea countries, and number of endemic species in each country.

Country	n. species mean ± s.d.	Range	Most common species (% localities)	n. endemic species
Algeria	3.38 ± 2.06	1-7	<i>Xenotrichula velox</i> (81%) <i>Paraturbanella teissieri</i> (62%)	1
Tunisia	1.00 ± 0.00	NA	NA NA	-
Malta	1.00 -	NA	NA NA	-
Egypt	12.86 ± 8.03	1-28	<i>Halichaetonotus spinosus</i> (64%) <i>Turbanella ambronerensis</i> (54%)	27
Israel	10.80 ± 5.16	2-17	<i>Halichaetonotus spinosus</i> (67%) <i>Acanthodasys aculeatus</i> (67%)	14
Cyprus	8.56 ± 5.36	1-17	<i>Draculiciteria tessellata</i> (33%) <i>Tetranchyroderma heterotubulatum</i> (67%)	7
Bulgaria	5.00 ± 3.00	2-	<i>Turbanella cornuta</i> (100%) <i>Halichaetonotus pleuracanthus</i> (67%)	-
Romania	7.57 ± 4.20	1-12	<i>Aspidiophorus mediterraneus</i> (43%) <i>Acanthodasys aculeatus</i> (43%)	1
Greece	8.24 ± 5.74	1-23	<i>Heteroxenotrichula squamosa</i> (39%) <i>Macrodasys caudatus</i> (50%)	12
Croatia	11.00 -	NA	NA NA	3
Italy	8.42 ± 6.96	1-37	<i>Halichaetonotus aculifer</i> (33%) <i>Acanthodasys aculeatus</i> (29%)	71
France	5.43 ± 5.43	1-26	<i>Paraturbanella dobrni</i> (24%) <i>Heteroxenotrichula squamosa</i> (19%)	5
Spain	3.00 -	Na	NA NA	-
Global data	8.04 ± 6.69	1-37	<i>Halichaetonotus aculifer</i> (28%) <i>Acanthodasys aculeatus</i> (28%)	141

NA, not applicable.

At species level, about 140 taxa have so far been found only in the Mediterranean; based on comparison with the well studied north European fauna (including the British Islands) it may be said that the endemicity rate of the basin is genuinely relevant. The remainder species (about 49% of total)




Fig. 4 - Representative species belonging to families and genera found in the Mediterranean basin. A-I: Thaumastodermatidae, A: *Hemidaysys agaso*; B: *Platydaysys phacellatus*; C: *Pseudostomella etrusca*; D: *Ptychostomella tyrrhenica*; E: *Tetranchyroderma cirrophorum*; F: *T. esarabdophorum*; G: *T. papii*; H: *T. thysanophorum*; I: *Thamastoderma ramuliferum*. L-N: Turbanellidae, L: *Paraturbanella pallida*; M: *Turbanella ambronensis*; N: *T. cornuta*.

have a geographic range extending to the North-Atlantic, and several (e.g., Fig. 5H) are considered regional cosmopolitans (i.e., reported from at least two Oceans). Within the Mediterranean, the species geographic distribution does not appear to be homogeneous, with several taxa restricted to either the western or to the Levantine basin, yet, Italy, Egypt and Israel show the highest numbers of endemic species, 70, 24 and 14, respectively. Of the widely distributed species, *Halichaetonotus atlanticus*, *Paraturbanella aggregotubulata* and *Pseudostomella cataphracta* seems restricted to the




Fig. 5 - Selected gastrotrich species found in the Mediterranean basin. A,B: The most common species of the two orders, A: the macrodasyidan *Acanthodaysys aculeatus*; B: the chaetonotidan *Halichaetonotus aculifer*. C,D: Species typical of the littoral zone, C: *Xenotrichula intermedia*; D: *Turbanella ambronensis*. E,F: Brackish-water species, E: *Heterolepidoderma foliatum*, F: *Dendrodaysys ponticum*. G,H: Species inhabiting organogenous sand, G: *Platydysys phacellatus*; H: *Diplodaysys ankeli*. A,E,G,H: SEM micrographs. B,C,D,F - DIC micrographs. Bar = A: 200 µm; B,C,E: 20 µm; D,F,G,H: 100 µm.

western basin, whereas *Dactylopoda baltica* and *Megadasys sterreri* are confined within the eastern area.

Only one species, *Chaetonotus antipai*, seems restricted to the Black Sea, testifying exchange of euryhaline species between the two Seas and/or the persistence of Sarmatic elements, which when tied to selected environments (i.e., brackish waters) were able to overcome the Messinian saline crisis (Sarà, 1985; Hsu, 1987; Bianchi and Morri, 2000); the recent findings of *Dendrodasys ponticus* (Fig. 5F) and *Turbanella cf. ponticus* also in the brackish water of the northern Adriatic Sea, perhaps, stands testimony of this (Tongiorgi et al., 1999). Among meiobenthic taxa, faunistic liaisons between the Northern Adriatic and the Pontic brackish areas were also highlighted by Ceccherelli and Mistri (1990) in a study analysing the ecological and zoogeographic association of brackish water harpacticoid copepods in the Mediterraenan and Black Seas. In the case of the gastrotrich-fauna, a cluster analysis using the Bray-Curtis presence-absence similarity value to examine the relationship between the investigated locations (organised prior analysis in 11 main areas) shows a clear separation between the Black Sea and the Mediterranean domain and within this the Algerian-Tunisian area is separated from the remaining 9 (Fig. 6). The latter 9 areas appear arranged in two main clusters, one clearly harmonious, including the southernmost Levantine regions (i.e., Egypt, Israel and Cyprus), whereas the other appears geographically somewhat less homogeneous, in that it contentiously unites regions of the northwestern basin, the Adriatic Sea and the two Greek sub-regions. Going into detail of this apparently contrasted cluster, the region comprising coastal France and Spain (Formentera), appears separated from the one containing, arranged into two distinct subsets, the three Italian and the two Greek areas. It is possible that the apparently lower faunistic similarity between coastal France and the western areas of Italy derives from discrepancies in the sampling effort made in two regions. It should however be emphasised that, in general, it is difficult to say if the scenarios depicted by the cluster analysis reflects real faunistic similarity among areas, or artefacts due to the different methodology used to gather faunistics data. For instance, information from Algeria and Tunisia derives from samples taken in the littoral zone only, whilst in Romania only the shallow sublittoral has so far been investigated. Since it has been shown that the littoral area hosts a much less diversified gastrotrich fauna (Todaro et al., 1995, 2002; Todaro and Rocha, 2003), yet selected species are typical of these sediments (Fig. 5C,D), it is therefore likely that gastrotrichs biodiversity in these countries is well underestimated, this statement is reinforced by considering also the very few locations sampled in these coastal areas, which is also true for several others (Table V). In turn, the unexplored portion of the biodiversity may unveil faunisitic similarity among areas not pinpointed so far.




Fig. 6 - Classification of Mediterranean and Black Sea areas, following cluster analysis on a species by location matrix after pooling data into 11 main regions.

CONCLUSION

The Mediterranean Sea and the Black Sea host a rich and diversified gastrotrich-fauna, including about half of the nearly 500 marine gastrotrich species known world-wide; moreover, all marine family and vast majority of marine genera have representatives in the Mediterranean basin, with two being endemic. It is likely that species richness of the basin will increase, as investigation will explore new promising habitat such as submarine caves (cfr. Todaro et al., 2001b), deep-sea sandy substrata or brackish environments.

As the Mediterranean apparently harbours somewhat between 4% and 18% of the world marine macroscopic species (Bianchi and Morri, 2000), it can be said that comparatively speaking the Gastrotricha represent one of the best known taxon in the basin. However, notwithstanding the extensive data set available, the paucity and/or absence of data from zoogeographical meaningful areas may hamper, more or less extensively, the meaning of any zoogeographical analysis. Information from Spain, Morocco and Turkey (both Mediterranean and Black Sea shores) is strongly needed to improve our current perception about diversity and distribution of the Mediterranean gastrotrich-fauna. In absence of such information, a more conservative use of the available records (e.g., analysing only information obtained by authors using a standardised methods) may also prove to be a task well worth tackling. Possibly in the light of the statement by Todaro (1992) and Todaro et al. (2001a) who, commenting on the distributional pattern of Italian species pointed out the importance of the edaphic factors in shaping local gastrotrich communities (Fig. 5E,G,H) in a basin, such as the Mediterranean, that lacks clear

barriers to dispersal of this meiofaunal group. Progress in molecular biology techniques and the falling cost of the consumables make nowadays feasible to carry out also in small-bodied organisms studies of population genetics, crucial to add information on the temporal scale; this in turn may help decoupling ecological factors and geological events, and consequently shed some light about origin and causes of the Mediterranean Gastrotricha biodiversity.

According to Hummon and Hummon (1995a) of the species found only in Red Sea or the Mediterranean, 20% are shared by both; molecular investigation could finally help distinguish also “modern migrant via the Suez canal, from the Tethyan relict persisting from an ancient, shared fauna”.

REFERENCES

- BALSAMO M., TODARO M.A. 1987 - *Aspidiophorus polystictos*, a new marine species (Gastrotricha, Chaetonotida) and its life cycle. Boll. Zool., 54: 147-153.
- BALSAMO M., TODARO M.A. 1988 - Life history traits of two chaetonotids (Gastrotricha) under different experimental conditions. Invertebr. Reprod. Dev., 14: 161-176.
- BALSAMO M., TONGIORGI, P. 1995 - Gastrotricha. In: A. Minelli, S. Ruffo, S. La Posta (eds.). Checklist delle specie della fauna d'Italia, Calderini, Bologna, fasc. 7: 1-11.
- BALSAMO M., FREGNI E., TONGIORGI P. 1994 - Marine and freshwater Gastrotricha from the Island of Montecristo (Tuscan Archipelago, Italy), with the description of new species. Boll. Zool., 61: 217-227.
- BALSAMO M., FREGNI E., TONGIORGI P. 1995 - Marine Gastrotricha from the coasts of Sardinia (Italy). Boll. Zool., 62: 273-286.
- BALSAMO M., FREGNI E., TONGIORGI P. 1996 - Marine Gastrotricha from Sicily with the description of a new species of *Chaetonotus*. Ital. J. Zool., 63: 173-183.
- BALSAMO M., TODARO M.A., TONGIORGI P. 1992 - Marine gastrotrichs from the Tuscan Archipelago (Tyrrhenian Sea): II. Chaetonotida, with description of three new species. Boll. Zool., 59: 487-498.
- BALSAMO M., FERRAGUTI M., GUIDI L., TODARO M. A., TONGIORGI P. 2002 - Reproductive system and spermatozoa of *Paraturbanella teissieri* (Gastrotricha, Macrodayida): implications for sperm transfer modality in Turbanellidae. Zoomorphology, 121: 235-241.
- BALSAMO M., HUMMON W.D., TODARO M.A., TONGIORGI P. 1997 - Italian marine Gastrotricha: IV. Four new species of Chaetonotida. Ital. J. Zool., 64: 83-89.
- BIANCHI C.N., MORRI C. 2000 - Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. Mar. Pollut. Bull., 40: 367-376.
- BOADEN P.J.S. 1965a - Two new interstitial Gastrotricha of the family Thaumastodermatidae. Pubbl. Stn. Zool. Napoli, 34: 219-225.
- BOADEN P.J.S. 1965b - Interstitial fauna from Porto Paone. Pubbl. Stn. Zool. Napoli, 34: 235-239.
- CECCHERELLI V.U., MISTRI M. 1990 - Ecological and zoogeographical study of some Mediterranean associations of brackish water harpacticoids. Boll. Zool., 57, 73-81.
- CLAPARÈDE E. 1867 - Miscellaneous zoologiques. III. Type d'un nouveau genre de gastrotriches. Ann. Sci. Nat. Zool., 8: 16-23.
- CLAUSEN C. 1965 - New interstitial species of the family Thaumastodermatidae (Gastrotricha Macrodasyoidea). Sarsia, 21: 23-36.
- COULL B.C. 1985 - Long-term variability of estuarine meiobenthos: an 11 year study. Mar. Ecol. Prog. Ser., 24: 205-218.
- DELAMARE DEBOUTTEVILLE C. 1953 - Monographie d'une espèce. L'écologie et la répartition du Mystacocaride *Derocheilocaris remanei* Delamare et Chappuis, en Méditerranée. Vie Milieu, 4: 321-380.
- DELAMARE DEBOUTTEVILLE C. 1954 - Eaux souterraines littorales de la côte catalane française (mise au point faunistique). Vie Milieu, 5: 408-451.

- DE ZIO S. 1965 - Distribuzione del mesopsammon in rapporto alla distanza dalla linea di riva e alla distanza da un corso di acqua salmastra. *Boll. Zool.*, **32**: 525-537.
- DE ZIO S., GRIMALDI P. 1964 - Analisi comparativa del mesopsammon di due spiagge Pugliesi in rapporto ad alcuni fattori ecologici. *Arch. Bot. Biogeogr. Ital.*, **9**: 357-367.
- EVANS W. A., TODARO M. A., HUMMON W. D. 1993 - Eutrophication and gastrotrich diversity in the Northern Adriatic Sea. *Mar. Pollut. Bull.*, **26**: 268-272.
- FAIENZA M. G., PIZZUTO A., GRIMALDI-DE ZIO S. 2000 - Ulteriori dati sulla fauna gastrotricologica del litorale adriatico pugliese. *Biol. Mar. Medit.*, **7**: 680-682.
- FAIENZA M. G., FREGNI E., TONGIORGI P., GRIMALDI-DE ZIO S. 1999 - Una faunula a gastrotrichi di detrito coralligeno e di una grotta sottomarina del Basso Adriatico. *Biol. Mar. Medit.*, **6**: 385-387.
- FERRAGUTI M., BALSAMO M. 1994 - Sperm morphology and anatomy of the genital organs in *Mesodasys laticaudatus* Remane, 1951 (Gastrotricha: Macrodasyida). *J. Submicrosc. Cytol. Pathol.*, **26**: 21-28.
- FERRAGUTI M., BALSAMO M. 1995 - Comparative spermatology of Gastrotricha. In: B.G.M. Jamieson, J. Ausio, J.L. Justine (a cura di), Advances in spermatozoal phylogeny and taxonomy. *Mém. Mus. Natl. Hist. Nat.*, **166**: 105-117.
- FERRAGUTI M., BALSAMO M., FREGNI E. 1995 - The spermatozoa of three species of Xenotrichulidae (Gastrotricha, Chaetonotida): the two "dunne Nebengeissein" of spermatozoa in *Heteroxenotrichula squamosa* are peculiar para-acrosomial bodies. *Zoomorphology*, **115**: 151-159.
- FIZE A. 1957 - Premières résultats des recoltes de microfaune des sables effectuees sur la côte Languedocienne. *Vie Milieu*, **8**: 377-381.
- FIZE A. 1963 - Contribution à l'étude de la microfaune des sables littoraux du Golfe d'Aigues-Mortes. *Vie Milieu*, **14**: 669-774.
- FREGNI E. 1998a - A practice in the extraction of gastrotrichs from marine and freshwater sediments. *Ital. J. Zool.*, **65**: 203-206.
- FREGNI E. 1998b - The spermatozoa of macrodasyids gastrotrichs: observations by scanning electron microscopy. *Invert. Reprod. Dev.*, **34**: 1-11.
- FREGNI E., TONGIORGI P., FAIENZA M.G. 1998 - Two new species of *Urodasys* (Gastroricha, Macrodasyidae) with cuticular styllet. *Ital. J. Zool.*, **65**: 377-380.
- FREGNI E., FAIENZA, M.G., GRIMALDI-DE ZIO S., TONGIORGI P., BALSAMO M. 1999 - Marine gastrotrichs from the Tremiti Archipelago in the southern Adriatic Sea, with the description of two new species of *Urodasys*. *Ital. J. Zool.*, **66**: 183-194.
- GERLACH S.A. 1953 - Gastrotrichen aus dem Küstengrundwasser des Mittelmeeres. *Zool. Anz.*, **150**: 203-211.
- GIERE O., ELEFTHERIOU A., MURISON D.J. 1988 - Abiotic factors. In: R. P. Higgins, H. Thiel (eds.), Introduction to the study of meiofauna. Washington DC, Smithsonian Institution Press: 61-78.
- GRÜNSPAN T. 1908 - Beiträge zur Systematik der Gastrotrichen. *Zool. Jahrb. (Syst.)*, **26**: 214-256.
- d'HONDRT J.L. 1974 - Contribution à l'étude de la microfaune interstitielle des plages de l'ouest Algérien. *Vie Milieu*, **23**: 227-241.
- HSU K.J., 1987 - The Mediterranean was a desert: a voyage of the Glomar Challenger. Princeton University Press, 197 pp.
- HUMMON W.D. 2001a - Global database for marine Gastrotricha on CD. Athens, Ohio University Zoological Collections.
- HUMMON W.D. 2001b - Marine Gastrotricha. Proc. 11th International Meiofauna Conference, Boston, US, 15-20 July 2001 (abs.).
- HUMMON W.D. 2001c - Global database: marine Gastrotricha. Proc. 11th International Meiofauna Conference, Boston, US, 15-20 July 2001 (abs.).
- HUMMON W.D., HUMMON M.R. 1992 - Marine Gastrotricha of Israel: Preliminary report. *Am. Zool.*, **32** (A): 115 (abs).
- HUMMON W.D., HUMMON M.R. 1995a - Marine Gastrotricha of eastern Mediterranean and Red seas. Proc. 9th International Meiofauna Conference, Perpignan, France 3-7 July 1995 (abs.).
- HUMMON W.D., HUMMON M.R. 1995b - Marine Gastrotricha of the eastern Mediterranean (EMED) and Red Seas (RED). *Am. Zool.*, **35**: 12A (abs 60).
- HUMMON W.D., ROILOU E. 1994 - Marine Gastrotricha of Greece: a preliminary report. *Biol. Gallo-Hell.*, **2**: 279-289.
- HUMMON W.D., BALSAMO M., TODARO M.A. 1992 - Italian marine Gastrotricha: I. Six new and one redescribed species of Chaetonotida. *Boll. Zool.*, **59**: 499-516.
- HUMMON W.D., HUMMON M.R., MOSTAFA H.M. 1994 - Marine Gastrotricha of Mediterranean Egypt. *Am. Zool.*, **34**: 10A (abs 54).
- HUMMON W.D., TODARO M.A., HUMMON M.R. 1990b - Marine Gastrotricha of the central Mediterranean Sea. *Am. Zool.*, **30**: 20A (abs 109).
- HUMMON W.D., TODARO M. A., TONGIORGI P. 1993 - Italian Marine Gastrotricha: II. One new genus and ten new species of Macrodasyida. *Boll. Zool.*, **60**: 109-127.
- HUMMON W.D., TODARO M. A., BALSAMO M., TONGIORGI P. 1990a. Effects of pollution on marine Gastrotricha in the northwestern Adriatic Sea. *Mar. Pollut. Bull.*, **21**: 241-243.

- HUMMON W.D., TODARO M. A., BALSAMO M., TONGIORGI P. 1996 - Italian marine Gastrotricha: III. Four new pentancrous species of the genus *Terranchyroderma* (Macrodasyida, Thaumastodermatidae). Ital. J. Zool., **63**: 73-79.
- HUMMON W.D., TODARO M. A., TONGIORGI P., BALSAMO M. 1998 - Italian marine Gastrotricha: V. Four new and one redescribed species of Macrodasyida in the Dactylopolidae and Thaumastodermatidae. Ital. J. Zool., **65**: 109-119.
- LUPORINI P., TONGIORGI P. 1972 - Observation au microscope à balayage de quelques gastrotriches chaetonotoïdes. Cah. Biol. Mar., **13**: 299-303.
- LUPORINI P., MAGAGNINI G., TONGIORGI P. 1971 - Contribution à la connaissance des gastrotriches des côtes de Toscane. Cah. Biol. Mar., **12**: 433-455.
- LUPORINI P., MAGAGNINI G., TONGIORGI P. 1973a - Gastrotrichi macrodasioïdei delle coste della Toscana. Pubbl. Stn. Zool. Napoli, **38** (1970): 267-288.
- LUPORINI P., MAGAGNINI G., TONGIORGI P. 1973b - Chaetonotoid gastrotrichs of the Tuscan Coast. Boll. Zool., **40**: 31-40.
- PAPI F. 1957 - Tre nuovi Gastrotrichi mediterranei. Pubbl. Str. Zool. Napoli, **30**: 176-183.
- REMANE A. 1927a - Neue Gastrotricha Macrodasyoidea. Zool. Jahrb. Abt. Syst. (Jena), **54**: 203-242.
- REMANE A. 1927b - Gastrotricha, In: G. Grimpé (ed.), Die Tierwelt der Nord- und Ostsee. Leipzig, Akademische Verlagsgesellschaft: 1-56.
- REMANE A. 1951 - *Mesodays*, ein neues Genus der Gastrotricha Macrodasyoidea aus der Kieler Bucht. Kieler Meeresforsch., **8**: 102-105.
- REMANE A. 1952 - Zwei neue *Turbanella*-Arten aus dem marinen Küstengrundwasser. Kieler Meeresforsch., **9**: 62-65.
- REMANE A. 1961 - *Neodays uchidai* nov. spec., eine zweite *Neodays*-Art (Gastrotricha Chaetonotoidea). Kieler Meeresforsch., **17**: 85-88.
- RENAUD-MORNANT J. 1968 - Presence du genre *Polymerurus* en milieu marin, description de deux espèces nouvelles (Gastrotricha, Chaetonotidae). Pubbl. Stn. Zool. Napoli, **36**: 141-151.
- RODEWALD L. 1938 - Beitrag zur Kenntnis der Systematik und Oekologie der Gastrotrichenfauna Dobrogea's (Rumaenien). Zool. Anz., **11**: 206-216.
- RUDESCU L. 1966 - Gastrotrichi voi pentra Marea Neagra (Sistematica, ecologia si zoogeografia grupului). Hidrobiologia (Bucharest), **7**: 103-130.
- RUDESCU L. 1967 - Gastrotricha. In: Fauna Republicii Socialiste România, **2** (3), 289 pp.
- SARÀ M., 1985 - Ecological factors and their biogeographic consequences in the Mediterranean ecosystems. In: M.M. Moraitou, M.M. Apostolopoulou, V. Kirtsis (eds.), Mediterranean marine ecosystems: New York and London, Plenum Press: 1-17.
- SCHROM H. 1966a - Gastrotrichen aus Feinsanden der Umgebung von Venedig. Boll. Mus. Civ. St. Nat. Venezia, **17**: 31-45.
- SCHROM H. 1966b - Verteilung einiger Gastrotrichen im oberen Eulitoral eines nordadriatischen Sandstrandes. Veroeff. Inst. Meeresforsch. Bremerhaven, Sonderb., **2**: 95-104.
- SCHROM H. 1968 - Zur Kenntnis des Mesopsammals der Nordadria. II. Die Sande in der Umgebung von Rovinj und ihre Faunenmerkmale. Thalassia Jugosl., **4**: 31-38.
- SCHROM H. 1972 - Nordadriatische Gastrotrichen. Helgol. Wiss. Meeres., **23**: 286-351.
- SWEDMARK B. 1956 - Étude de la microfaune des sables marins de la région de Marseille. Arch. Zool. Exp. Gén., **93**: 70-95.
- TODARO M.A. 1992 - Contribution to the study of the Mediterranean meiofauna: Gastrotricha from the Island of Ponza, Italy. Boll. Zool., **59**: 321-333.
- TODARO M.A. 1997 - Meiofauna, il popolo della sabbia: Il microscopico mondo animale delle secche della Meloria. Livorno, Otello Debatté editore, 79 pp.
- TODARO M.A. 1998a - Meiofauna from the Meloria shoals: Gastrotricha biodiversity and seasonal dynamics. Biol. Mar. Medit., **5**: 585-590.
- TODARO M.A. 1998b - Note faunistiche ed ecologiche sui Gastrotrichi delle Secche della Meloria. In: Fauna nel Mediterraneo, immagini e note di ecologia marina. Quaderni dell'Acquario (Livorno), Tipografia Marengo, Livorno, vol. 3: 57-72.
- TODARO M.A., BALSAMO M. 1994 - Marine gastrotrichs from Sicily (Italy). Biol. Gallo-Hell., **22**: 291-292.
- TODARO M.A., ROCHA C.E.F. 2003 - Diversity and distribution of marine Gastrotricha along the northern beaches of the State of São Paulo (Brazil), with a description of *Macrodays fornerise* sp. nov. (Macrodasyida, Macrodasyidae). J. Nat. Hist. (in press).
- TODARO M.A., BALSAMO M., TONGIORGI P. 1988 - *Terranchyroderma sardum*, a new species of the family Thaumastodermatidae (Gastrotricha, Macrodasyida). Boll. Zool., **55**: 69-72.
- TODARO M.A., BALSAMO M., TONGIORGI P. 1992 - Marine gastrotrichs from the Tuscan Archipelago (Tyrrenian Sea): I. Macrodasyida, with description of three new species. Boll. Zool., **59**: 471-485.
- TODARO M.A., BALSAMO M., TONGIORGI P. 2002 - Marine gastrotrich fauna in Corsica (France), with description of a new species of *Terranchyroderma* (Gastrotricha, Macrodasyida). Mar. Biol. Assoc. U.K., **82**: 555-564.
- TODARO M.A., BALSAMO M., MOSCI D., TONGIORGI P. 2001b - Emergenze meiobentoniche di una grotta marina del litorale salentino (LE). Atti 62° congresso UZI, Sanremo 23-27 settembre.

- TODARO M.A., FLEEGER J.W., HUMMON W.D. 1995. Marine gastrotrichs from the sand beaches of the northern Gulf of Mexico: species list and distribution. *Hydrobiologia*, **310**: 107-117.
- TODARO M.A., FLEEGER J.W., HU Y.P., HRINCEVICH A.W., FOLTZ D.W. 1996 - Are meiofauna species cosmopolitan? Morphological and molecular analysis of *Xenotrichula intermedia* (Gastrotricha: Chaetonotida). *Mar. Biol.*, **125**: 735-742.
- TODARO M.A., HUMMON W.D., BALSAMO M., FREGNI E., TONGIORGI P. 2001a - Inventario dei gastrotrichi marini italiani: una checklist annotata. *Atti Soc. Tosc. Sci. Nat., Mem., Ser. B*, **107**: 75-137.
- TODARO M.A., LITTLEWOOD D.T.J., BALSAMO M., HERNIOU E., CASSANELLI S., MANICARDI G.C., WIRZ A., TONGIORGI P. 2003 - The interrelationships of the Gastrotricha using nuclear small rRNA subunit sequence data, with an interpretation based on morphology. *Zool. Anz.*, **242**: 145-156.
- TONGIORGI P. 1975 - Two interesting Macro dasy oidea (Gastrotricha) from the coast of Tuscany. *Boll. Zool.*, **42**: 275-278.
- TONGIORGI P., BALSAMO M. 1984 - A new *Tetranchyroderma* species (Gastrotricha, Macro dasy oidea) from the Adriatic coast. *Boll. Zool.*, **51**: 335-338.
- TONGIORGI P., FREGNI E., BALSAMO M. 1999 - Gastrotricha from Italian brackish environment with description of a new species of *Chaetonotus*. *J. Mar. Biol. Ass. U.K.*, **79**: 585-592.
- VALKANOV A. 1957 - Erster Versuch zur erforschung der Gastrotrichen des Schwarzen Meeres. *Ann. Univ. Sofia Fac. Biol.*, **50**: 383-399.
- VIVIER M.H. 1974 - *Musellifer profundus* n. sp. Gastrotriche (Chaetonotidae) des vases profondes des Mediterranee. *Bull. Soc. Zool. France*, **99**: 183-186.
- WESTHEIDE W. 1972 - Räumliche und zeitliche Differenzierungen im Verteilungsmuster der marinen Interstitialfauna. *Deutsch. Zool. Ges. Ver.*, **65**: 23-32.
- WILKE U. 1954 - Mediterrane Gastrotrichen. *Zool. Jahrb. Abt. Syst. (Jena)*, **82**: 497-550.
- WIRZ A., PUCCIARELLI S., MICELI C., TONGIORGI P., BALSAMO M. 1999 - Novelty in phylogeny of Gastrotricha: evidence from 18S rRNA gene. *Mol. Phylog. Evol.*, **13**: 314-318.

Appendix I – Species list of Gastrotricha Macrodayida found so far in 11 Mediterranean and 2 Black Sea countries.

Taxon

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Romania	Greece	Croatia	Italy	Malta	France	Spain
Family DACTYLOPODOLIDAE													
Genus <i>Dactylopodola</i> Strand, 1929													
<i>Dactylopodola baltica</i> (Remane, 1926)	-	-	+	-	-	-	-	+	-	-	-	-	-
<i>Dactylopodola mesotyphle</i> Hummon et al., 1998	-	-	+	-	+	-	-	+	-	+	-	+	-
<i>Dactylopodola typhle</i> (Remane, 1927)	+	-	+	+	+	-	+	-	-	+	-	+	-
Genus <i>Dendrodaysys</i> Wilke, 1954													
<i>Dendrodaysys affinis</i> Wilke, 1954	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Dendrodaysys gracilis</i> Wilke, 1954	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Dendrodaysys ponticus</i> Valkanov, 1957	-	-	-	-	-	+	+	-	-	+	-	-	-
Genus <i>Dendropodola</i> Hummon et al., 1993													
<i>Dendropodola transitionalis</i> Hummon et al., 1993	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Xenodaysys</i> Swedmark, 1967													
<i>Xenodaysys</i> sp. A [Todaro et al., 2003]	-	-	-	-	-	-	-	-	-	+	-	-	-
Family LEPIDODASYIDAE													
Genus <i>Cephalodaysys</i> Remane, 1926													
<i>Cephalodaysys hadrosomus</i> Hummon et al., 1993	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Cephalodaysys littoralis</i> Renaud-Debyser, 1964	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Cephalodaysys palavensis</i> Fize, 1963	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>Cephalodaysys swedmarki</i> **	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>Cephalodaysys turbanelloides</i> (Boaden, 1960)	-	-	-	-	-	-	-	+	-	+	-	+	-
<i>Cephalodaysys</i> sp. A Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
Genus <i>Dolichodaysys</i> Gagne, 1977													
<i>Dolichodaysys elongatus</i> Gagne, 1977	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Lepidodaysys</i> Remane, 1926													
<i>Lepidodaysys martini</i> Remane, 1926	-	-	-	-	-	-	-	-	-	+	-	+	-
<i>Lepidodaysys platyurus</i> Remane, 1927	-	-	-	-	-	-	-	+	-	-	+	-	-
<i>Lepidodaysys unicarenatus</i> Balsamo et al., 1994	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Lepidodaysys</i> sp. A [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Megadaysys</i> Schmidt, 1974													
<i>Megadaysys minor</i> Ksielewski, 1987	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Megadaysys sterreri</i> (Boaden, 1974)	-	-	+	-	+	-	-	-	-	-	-	-	-
Genus <i>Mesodaysys</i> Remane, 1951													
<i>Mesodaysys adenotubulatus</i> Hummon et al., 1993	-	-	+	+	-	-	-	+	-	+	-	+	+
<i>Mesodaysys ischiensis</i> Hummon et al., 1993	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Mesodaysys laticaudatus</i> Remane, 1951	-	-	+	+	-	-	-	+	-	+	-	+	-
<i>Mesodaysys littoralis</i> Remane, 1951	+	-	+	+	+	-	-	+	-	+	-	+	-
<i>Mesodaysys</i> sp. A Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
Genus <i>Paradaysys</i> Remane, 1934													
<i>Paradaysys subterraneus</i> Remane, 1934	-	-	-	-	-	-	-	-	-	+	-	+	-
<i>Paradaysys</i> sp. A France*	-	-	-	-	-	-	-	-	-	-	-	+	-
Genus <i>Pleurodaysys</i> Remane, 1927													
<i>Pleurodaysys helgolandicus</i> Remane, 1927	-	-	-	-	-	-	-	-	-	+	-	+	-
Family MACRODASYIDAE													
Genus <i>Macrodaysys</i> Remane, 1924													

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Romania	Greece	Croatia	Italy	Malta	France	Spain
<i>Macrodasy s africanus</i> Valkanov, 1957	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s balticus</i> Roszczak, 1939	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s buddenbrocki</i> Remane, 1924	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s caudatus</i> Remane, 1927	-	-	+	+	-	-	-	-	-	-	-	-	-
<i>Macrodasy s gerlachi</i> Papi, 1957	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s neapolitanus</i> Papi, 1957	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s thucus</i> Laporini et al., 1973	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. A Cyprus*	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. E Greece*	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. A Greece*	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. A Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. B Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. C Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. D Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. J Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. K Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. M Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. N Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. S Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. C Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. D Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. E Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Macrodasy s</i> sp. B [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Macrodasy s</i> sp. C [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Macrodasy s</i> sp. E [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	-	+	-	-
Genus <i>Uroday s</i> Remane, 1926													
<i>Uroday s acanthostylis</i> Fregni et al., 1998	-	-	-	-	-	-	-	-	-	-	+	+	-
<i>Uroday s apulensis</i> Fregni et al., 1999	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Uroday s bucinastylis</i> Fregni et al., 1999	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Uroday s viviparus</i> Wilke, 1954	-	-	-	-	+	-	-	-	-	-	+	-	-
Family PLANODASYIDAE													
Genus <i>Crasiella</i> Clausen, 1968													
<i>Crasiella</i> sp. A [Fregni et al., 1999]	-	-	-	-	-	-	-	-	-	-	+	-	-
Family THAUMASTODERMATIDAE													
Genus <i>Acanthodasy s</i> Remane, 1927													
<i>Acanthodasy s aculeatus</i> Remane, 1927	-	-	+	+	+	+	+	+	-	+	-	+	-
<i>Acanthodasy s</i> sp. A [Schrom, 1972]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Acanthodasy s</i> sp. B [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Acanthodasy s</i> sp. C [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Diplodasy s</i> Remane, 1927													
<i>Diplodasy s ankeli</i> Wilke, 1954	-	-	+	-	-	-	-	-	-	+	-	+	-
<i>Diplodasy s meloriae</i> Todaro et al., 1992	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Diplodasy s minor</i> Remane, 1926	-	-	-	-	-	-	+	+	-	+	-	-	-
<i>Diplodasy s platydayooides</i> Remane, 1927	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Diplodasy s</i> sp. A Cyprus*	-	-	-	-	+	-	-	-	-	-	-	-	-

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Romania	Greece	Croatia	Italy	Malta	France	Spain
<i>Diplodasy</i> sp. A [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Hemidasy</i> Claparède, 1867	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Hemidasy agaso</i> Claparède, 1867	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Platydasy</i> Remane, 1927	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Platydasy maximus</i> Remane, 1927	-	-	-	-	-	-	-	+	-	-	+	-	-
<i>Platydasy ocellatus</i> Clausen, 1965	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Platydasy phacellatus</i> Clausen, 1965	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Platydasy ruber</i> Swedmark, 1956	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Platydasy stiliferus</i> Boaden, 1965	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Platydasy tentaculatus</i> Swedmark, 1956	-	-	-	-	-	-	-	-	-	-	+	-	-
Genus <i>Pseudostomella</i> Swedmark, 1956	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Pseudostomella cataphracta</i> Ruppert, 1970	-	-	-	-	-	-	-	-	-	-	+	-	-
<i>Pseudostomella etrusca</i> Hummon et al., 1993	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Pseudostomella roscovita</i> Swedmark, 1956	+	-	-	-	-	-	-	+	-	+	-	+	-
Genus <i>Ptychostomella</i> Remane, 1926	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Ptychostomella mediterranea</i> Remane, 1927	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Ptychostomella tyrrhenica</i> Hummon et al., 1993	-	-	+	-	+	-	-	-	-	+	-	+	-
Genus <i>Tetranchyroderma</i> Remane, 1926	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tetranchyroderma anomalopsum</i> Humm. et al., 1996	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma antennatum</i> Luporini et al., 1973	-	+	-	-	-	-	-	+	-	+	-	-	-
<i>Tetranchyroderma aphenothignum</i> Humm. et al., 1998	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Tetranchyroderma apum</i> Remane, 1927	-	-	-	-	-	-	+	-	-	+	-	-	-
<i>Tetranchyroderma boadeni</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma cirrophorum</i> Levi, 1950	-	-	-	+	-	-	-	-	-	+	-	+	-
<i>Tetranchyroderma colopodium</i> Boaden, 1963	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma esarabdophorum</i> Tong, et al., 1984	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Tetranchyroderma heterotubulatum</i> Humm. et al., 1993	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Tetranchyroderma hirtum</i> Luporini et al., 1973	-	-	-	-	-	-	-	-	-	+	-	+	-
<i>Tetranchyroderma hypopsilancrum</i> Humm. et al., 1993	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma inaequitubulatum</i> Todaro et al., 2002	-	-	-	-	-	-	-	-	-	+	-	+	-
<i>Tetranchyroderma insulare</i> Balsamo et al., 1994	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma kontosorum</i> Hummon et al., 1996	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma masilense</i> Swedmark, 1956	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>Tetranchyroderma megastomum</i> (Remane, 1927)	-	-	+	-	-	-	-	+	-	+	-	+	-
<i>Tetranchyroderma pachysomum</i> Hummon et al., 1993	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma papii</i> Gerlach, 1953	+	+	+	+	+	-	-	+	-	+	-	+	-
<i>Tetranchyroderma polypodium</i> Luporini et al., 1973	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma polypyrobolostomum</i> Humm. et al., 1996	-	-	+	-	-	-	-	+	-	+	-	-	-
<i>Tetranchyroderma psilotopum</i> Hummon et al., 1998	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma quadridentaculatum</i> Tod. et al., 1992	-	-	-	-	-	-	-	+	-	+	-	+	-
<i>Tetranchyroderma sanctaecaterinae</i> Todaro et al., 1992	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma sardum</i> Todaro et al., 1988	-	-	-	-	-	-	-	+	-	+	-	+	-
<i>Tetranchyroderma symphorochetum</i> Hummon et al., 1998	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma tanymesatherum</i> Hummon et al., 1996	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranchyroderma ibisanogaster</i> Boaden, 1965	-	-	-	-	-	-	-	-	-	+	-	-	-

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Romania	Greece	Croatia	Italy	Malta	France	Spain
<i>Tetranychoderma thysanophorum</i> Hummon et al., 1993	-	-	-	-	-	-	-	+	-	+	-	+	-
<i>Tetranychoderma verum</i> Wilke, 1954	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Tetranychoderma</i> sp. A Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Tetranychoderma</i> sp. B Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Tetranychoderma</i> sp. A Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Tetranychoderma</i> sp. D Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Tetranychoderma</i> sp. A [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. B [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. AB [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. AC [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. AD [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. W [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. II [Schrom, 1972]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Tetranychoderma</i> sp. III [Schrom, 1972]	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Thaumastoderma</i> Remane, 1927													
<i>Thaumastoderma bifurcatum</i> Clausen, 1991	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Thaumastoderma heideri</i> Remane, 1926	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Thaumastoderma mediterraneum</i> Remane, 1927	-	-	-	-	-	-	-	+	+	+	-	+	-
<i>Thaumastoderma ramuliferum</i> Clausen, 1965	-	-	-	-	+	-	-	-	-	+	-	+	-
<i>Thaumastoderma swedmarki</i> Levi, 1959	-	-	-	-	-	-	-	-	-	-	-	+	-
Family TURBANELLIDAE													
Genus <i>Paraturbanella</i> Remane, 1927													
<i>Paraturbanella aggregotubulata</i> Evans, 1992	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>Paraturbanella dobrni</i> Remane, 1927	-	-	-	-	-	-	-	+	-	+	-	+	-
<i>Paraturbanella levantina</i> **	+	-	-	-	-	-	-	-	-	-	-	-	-
<i>Paraturbanella pallida</i> Luporini et al., 1973	-	-	+	-	-	-	-	+	-	+	-	+	-
<i>Paraturbanella teissieri</i> Swedmark, 1954	+	-	+	-	-	-	-	+	-	+	-	+	+
<i>Paraturbanella</i> sp. A Egypt*	-	-	+	-	+	-	-	-	-	-	-	-	-
Genus <i>Turbanella</i> Schultze, 1853													
<i>Turbanella ambronensis</i> Remane, 1943	+	-	+	+	+	-	-	+	-	+	-	+	-
<i>Turbanella bocqueti</i> Kaplan, 1958	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Turbanella cornuta</i> Remane, 1924	+	-	+	-	-	+	+	+	-	+	-	+	-
<i>Turbanella hyalina</i> Schultze, 1853	-	-	-	-	-	-	+	+	-	+	-	+	-
<i>Turbanella otti</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Turbanella petiti</i> Remane, 1952	-	-	-	-	-	-	-	-	-	+	-	+	-
<i>Turbanella pontica</i> Valkanov, 1957	-	-	-	-	-	+	+	-	-	+	-	-	-
<i>Turbanella subterranea</i> Remane, 1935	-	-	-	+	-	-	+	-	-	+	-	-	-
<i>Turbanella veneziana</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Turbanella</i> sp. A Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-

*Undescribed species, reported in Hummon (2001a).

***Nomen nudum*, reported in Hummon (2001a).

Appendix II – Species list of Gastrotricha Chaetonotida found so far in 11 Mediterranean and 2 Black sea countries.

Taxon

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Greece	Croatia	Italy	Malta	France	Spain	Romania
Suborder MULTITUBULATINA													
Family NEODASYIDAE													
Genus <i>Neodasys</i> Remane, 1927													
<i>Neodasys uchidai</i> Remane, 1961	-	-	-	-	-	-	-	-	-	-	-	+	-
Suborder PAUCITUBULATINA													
Family XENOTRICHULIDAE													
Genus <i>Draculiciteria</i> Hummon, 1974													
<i>Draculiciteria tesserata</i> (Renaud-Mornant, 1968)	-	-	+	+	+	-	-	+	-	+	-	+	-
Genus <i>Heteroxenotrichula</i> Wilke, 1954													
<i>Heteroxenotrichula affinis</i> (Remane, 1934)	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Heteroxenotrichula arcasonensis</i> Ruppert, 1979	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Heteroxenotrichula pygmaea</i> (Remane, 1934)	-	-	+	+	-	+	-	+	-	+	-	+	-
<i>Heteroxenotrichula pseudosquamosa</i> **	-	-	+	+	-	-	-	-	-	-	-	-	-
<i>Heteroxenotrichula squamosa</i> Wilke, 1954	-	-	+	+	+	-	-	+	-	+	-	+	+
<i>Heteroxenotrichula subterranea</i> (Remane, 1934)	-	-	+	+	-	-	-	-	-	+	-	-	-
<i>Heteroxenotrichula</i> sp. A Egypt*	-	-	+	+	-	-	-	-	-	-	-	-	-
Genus <i>Xenotrichula</i> Remane, 1927													
<i>Xenotrichula bispina</i> Roszczak, 1957	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Xenotrichula cornuta</i> Wilke, 1954	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Xenotrichula intermedia</i> Remane, 1934	+	-	+	+	+	+	+	+	+	+	-	+	-
<i>Xenotrichula lineata</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Xenotrichula punctata</i> Wilke, 1954	-	-	+	-	-	-	-	+	-	+	-	+	-
<i>Xenotrichula soikai</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Xenotrichula velox</i> Remane, 1927	+	-	-	-	-	-	-	+	-	-	-	+	-
Family CHAETONOTIDAE													
Genus <i>Aspidiophorus</i> Voigt, 1904													
<i>Aspidiophorus lamellophorus</i> Hummon et al., 1992	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Aspidiophorus marinus</i> Remane, 1926	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Aspidiophorus mediterraneus</i> Remane, 1927	-	-	+	-	-	+	+	+	+	+	-	+	-
<i>Aspidiophorus paramediterraneus</i> Hummon, 1974	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Aspidiophorus polystictos</i> Balsamo & Todaro, 1987	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Aspidiophorus tentaculatus</i> Wilke, 1954	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Aspidiophorus</i> sp. Israel A*	-	-	-	+	-	-	-	-	-	-	-	-	-
Genus <i>Chaetonotus</i> Eherenberg, 1830													
Subgenus <i>Euchaetonotus</i> Schwank, 1990													
<i>Chaetonotus aegilonensis</i> Balsamo et al., 1992	-	-	+	-	+	-	-	+	-	+	-	+	-
<i>Chaetonotus aequispinosus</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonotus angustus</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonotus antipai</i> Rodewald, 1938	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Chaetonotus apecochaeetus</i> Hummon et al., 1992	-	-	+	-	+	-	-	+	-	+	-	+	-

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Greece	Croatia	Italy	Malta	France	Spain	Romania
<i>Chaetonus apolemmus</i> Hummon et al., 1992	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Chaetonus ichthyoidoides</i> Tongiorgi et al., 1999	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus magnificus</i> Balsamo et al., 1997	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus mariae</i> Todaro, 1992	-	-	+	-	+	-	-	-	-	-	-	-	-
<i>Chaetonus maximus</i> Ehrenberg, 1830	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Chaetonus mediterraneus</i> Balsamo et al., 1997	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Chaetonus napoleonicus</i> Balsamo et al., 1992	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonus parthenopeus</i> Wilke, 1954	+	-	-	+	-	-	-	-	-	+	-	-	-
<i>Chaetonus siciliensis</i> Hummon et al., 1992	-	-	-	-	-	-	-	+	-	+	-	+	-
<i>Chaetonus similis</i> Zelinka, 1889	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Chaetonus</i> sp. E [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonus</i> sp. F [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonus</i> sp. P [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonus</i> sp. Q [Todaro et al., 2001]	-	-	+	+	+	-	-	+	-	-	-	+	-
Subgenus <i>Hystricochaetonotus</i> Schwank, 1990													
<i>Chaetonus lacunosus</i> Mock, 1979	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Chaetonus variosquamatus</i> Mock, 1979	-	-	+	+	-	-	-	+	-	+	-	+	-
Subgenus <i>Schizochaetonotus</i> Schwank, 1990													
<i>Chaetonus atrox</i> Wilke, 1954	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Chaetonus dispar</i> Wilke, 1954	-	-	+	+	-	-	-	+	+	+	-	+	-
<i>Chaetonus hilaris</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonus inaequidentatus</i> Kisielewski, 1988	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Chaetonus jueundus</i> (d'Hondt, 1971)	-	-	+	+	-	-	-	-	-	-	-	-	-
<i>Chaetonus luporinii</i> Balsamo et al., 1996	-	-	-	-	+	-	-	-	-	+	-	-	-
<i>Chaetonus modestus</i> Schrom, 1972	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chaetonus neptuni</i> Wilke, 1954	-	-	+	-	+	-	-	+	-	+	-	+	-
<i>Chaetonus schromi</i> Hummon, 1974	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chaetonus serenus</i> Schrom, 1972	-	-	-	-	-	-	-	-	+	+	-	-	-
<i>Chaetonus balticus</i> Remane, 1926	-	-	-	-	-	-	+	-	-	-	-	-	-
Subgenus unknown													
<i>Chaetonus</i> sp. A Cyprus*	-	-	+	-	+	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. B Cyprus*	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. C Cyprus*	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. D Cyprus*	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. E Cyprus*	-	-	-	-	+	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. A Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. B Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. C Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. F Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. G Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. H Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonus</i> sp. J Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Greece	Croatia	Italy	Malta	France	Spain	Romania
<i>Chaetonotus</i> sp. K Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonotus</i> sp. L Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonotus</i> sp. M Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Chaetonotus</i> sp. E England*	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Chaetonotus</i> sp. A Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Chaetonotus</i> sp. C Greece*	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chaetonotus</i> sp. H Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Chaetonotus</i> sp. A Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Chaetonotus</i> sp. F Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
Genus <i>Halichaeonotus</i> (Remane, 1936)													
<i>Halichaeonotus aculifer</i> (Gerlach, 1953)	+	-	+	+	+	-	+	+	-	+	-	+	-
<i>Halichaeonotus atlanticus</i> Kisielewski, 1988	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus batillifer</i> (Luporini et al., 1972)	-	-	-	+	-	-	-	+	-	+	-	-	-
<i>Halichaeonotus clavicornis</i> Balsamo et al., 1995	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus decipiens</i> (Remane, 1926)	-	-	+	-	-	+	+	+	-	+	-	+	-
<i>Halichaeonotus etroloimus</i> Hummon et al., 1992	-	-	+	-	-	-	+	-	+	-	-	-	-
<i>Halichaeonotus genatus</i> Balsamo et al., 1995	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus italicus</i> Balsamo, et al., 1997	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus jucundus</i> (d'Hondt, 1971)	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus margaretae</i> Hummon et al., 1992	-	-	+	+	-	-	-	+	-	+	-	-	-
<i>Halichaeonotus marivagus</i> Balsamo et al., 1992	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Halichaeonotus paradoxus</i> (Remane, 1927)	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Halichaeonotus parvus</i> (Wilke, 1954)	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus pleuracanthus</i> (Remane, 1927)	-	-	-	-	+	+	-	-	-	-	-	+	-
<i>Halichaeonotus riedli</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus spinosus</i> Mock, 1979	-	-	+	+	+	-	-	+	-	+	-	+	-
<i>Halichaeonotus swedmarki</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus thalassopais</i> Hummon et al., 1992	-	-	+	+	-	-	-	+	-	+	-	+	-
<i>Halichaeonotus</i> sp. A Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Halichaeonotus</i> sp. A Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Halichaeonotus</i> sp. B Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Halichaeonotus</i> sp. G Greece*	-	-	-	-	-	-	-	+	-	-	-	-	-
<i>Halichaeonotus</i> sp. B Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Halichaeonotus</i> sp. V [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Halichaeonotus</i> sp. Z [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Heterolepidoderma</i> Remane, 1926													
<i>Heterolepidoderma armatum</i> Schrom, 1966	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Heterolepidoderma clipeatum</i> Schrom, 1972	-	-	-	+	-	-	-	+	+	-	-	-	-
<i>Heterolepidoderma connectum</i> Schrom, 1972	-	-	-	+	-	-	-	+	+	-	-	-	-
<i>Heterolepidoderma foliatum</i> Renaud-Mornant, 1967	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Heterolepidoderma hermafroditum</i> Wilke, 1954	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Heterolepidoderma istriatum</i> Schrom, 1972	-	-	-	-	+	-	-	+	+	-	-	-	-

	Algeria	Tunisia	Egypt	Israel	Cyprus	Bulgaria	Greece	Croatia	Italy	Malta	France	Spain	Romania
<i>Heterolepidoderma loricatum</i> Schrom, 1972	-	-	+	+	+	-	-	+	-	+	-	-	-
<i>Heterolepidoderma marinum</i> Remane, 1926	-	-	-	-	-	+	+	-	-	-	-	-	-
<i>Heterolepidoderma ocellatum</i> (Metschnikoff, 19864)	-	-	-	-	-	+	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. A Cyprus*	-	-	+	-	+	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. A Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. B Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. D Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. I Egypt*	-	-	+	-	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. A Greece*	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. B Greece*	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. C Greece*	-	-	-	-	-	-	+	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. A Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. B Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. D Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. E Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. F Israel*	-	-	-	+	-	-	-	-	-	-	-	-	-
<i>Heterolepidoderma</i> sp. A [Todaro et al., 2001]	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>Heterolepidoderma</i> sp. B [Schrom, 1972]	-	-	-	-	-	-	-	+	-	+	-	-	-
<i>Heterolepidoderma</i> sp. C [Todaro et al., 2001]	-	-	-	-	-	-	-	-	-	+	-	-	-
Genus <i>Ichthydium</i> Ehrenberg, 1830													
<i>Ichthydium podura</i> (O.F. Müller, 1773)	-	-	-	-	-	+	+	-	+	-	-	-	-
<i>Ichthydium tergestinum</i> Gruenspan, 1908	-	-	-	-	-	+	-	-	+	-	-	-	-
Genus <i>Lepidodermella</i> Blake, 1933													
<i>Lepidodermella limogena</i> Schrom, 1972	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Lepidodermella squamata</i> (Dujardin, 1841)	-	-	-	-	-	+	-	-	+	-	-	-	-
Genus <i>Musellifer</i> Hummon, 1969													
<i>Musellifer delamarei</i> (Renaud-Mornant, 1968)	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>Musellifer profundus</i> Vivier, 1974	-	-	-	-	-	-	-	-	-	-	-	+	-

*Undescribed species, reported in Hummon (2001a).

***Nomen nudum*, reported in Hummon (2001a).