

# Lawrence Berkeley National Laboratory

## Lawrence Berkeley National Laboratory

### Title

THE ISOTOPES OF AMERICIUM

### Permalink

<https://escholarship.org/uc/item/95g2s4dd>

### Authors

Street, K.  
Ghirosio, A.  
Seaborg, G.T.

### Publication Date

2008-06-25

UCRL 655  
Copy ~~6972~~

C.2

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*


TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-655  
C2





UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

*DECLASSIFIED*

THE ISOTOPES OF AMERICIUM

K. Street, Jr., A. Ghiorso, and G. T. Seaborg

April 11, 1950

CAUTION

This document contains information affecting the National Defense of the United States. Its transmission or the disclosure of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalties under applicable Federal laws.

Berkeley, California

DECLASSIFIED

UCRL-655

Page 2

Chemistry-Transuranic Elements

STANDARD DISTRIBUTION

No. of Copies

Argonne National Laboratory	1-10
Atomic Energy Commission, Washington	11-12
Brookhaven National Laboratory	13-16
Carbide and Carbon Chemicals Corporation (K-25)	17-20
General Electric Company, Richland	21-26
Hanford Operations Office	27
Iowa State College	28
Knolls Atomic Power Laboratory	29-32
Los Alamos	33-35
Mound Laboratory	36-38
Naval Radiological Defense Laboratory	39
NEPA Project	40
New York Operations Office	41-42
Oak Ridge National Laboratory	43-50
Patent Branch, Washington	51
Technical Information Branch ORE	52-66
UCLA Medical Research Laboratory (Warren)	67
University of California Radiation Laboratory	68-70
University of Rochester	71-72

INFORMATION DIVISION  
Radiation Laboratory  
Univ. of California  
Berkeley, California

## THE ISOTOPES OF AMERICIUM

K. Street, Jr., A. Ghiorso, and G. T. Seaborg  
Radiation Laboratory and Department of Chemistry  
University of California, Berkeley, California

April 11, 1950

## ABSTRACT

Three new americium activities ( $\text{Am}^{238?}$ ,  $\text{Am}^{243}$ , and  $\text{Am}^{244?}$ , the latter two formed by  $n, \gamma$  reactions) are described and some additional information is given on previously reported americium isotopes.

DECLASSIFIED

THE ISOTOPES OF AMERICIUM

K. Street, Jr., A. Ghiorso, and G. T. Seaborg  
Radiation Laboratory and Department of Chemistry  
University of California, Berkeley, California  
April 11, 1950

Intermittent investigation of the isotopes of americium during the past two years has resulted in the production of three new americium activities and has also yielded some additional information about previously<sup>1</sup> reported activities. Since the work on americium is still in progress, this letter will give only a brief report of the new results obtained and a detailed description of the experiments will be deferred until a later date.

Am<sup>238?</sup>.-- Bombardment of Pu<sup>239</sup> with 50-Mev deuterons in the 184-inch cyclotron results in the production of a new americium activity of ca. 1.2 hours half-life in addition to the previously reported 12- and 50-hour activities.<sup>1</sup> Differential counting with beryllium and lead absorbers indicate that the decay is accompanied by the emission of conversion electrons and x-rays. The same activity is produced in good yield with 17-Mev deuterons in the 60-inch cyclotron, thus its mass number is probably greater than 237, since there is good evidence<sup>2,3</sup> that even with the heavy elements the yield of the (d,4n) reaction is very small at 17 Mev. In view of its half-life, radiation characteristics, and method of formation, this activity is probably an orbital electron capturing isotope and is best assigned to Am<sup>238</sup>.

---

<sup>1</sup>G. T. Seaborg, R. A. James, and L. O. Morgan, National Nuclear Energy Series, Plutonium Project Record, Vol. 14B, "The Transuranium Elements: Research Papers," Paper 22.2 (McGraw-Hill Book Co., Inc., New York, 1949).

<sup>2</sup>R. A. James, A. E. Florin, H. H. Hopkins, Jr., and A. Ghiorso, ibid., Paper No. 22.8.

<sup>3</sup>L. B. Magnusson, S. G. Thompson, and G. T. Seaborg, Phys. Rev. (May 15, 1950 issue).

Am<sup>239</sup>.-- The 12-hour orbital electron capturing Am<sup>239</sup> has been previously reported to decay by alpha-particle emission of unknown energy to the extent of ca. 0.1 percent.<sup>1</sup> A redetermination of the alpha branching ratio yielded the better value ca. 0.01 percent. The energy of the alpha-particle has been determined by means of a multi-channel pulse analyzer to be  $5.77 \pm 0.05$  Mev.

Am<sup>240</sup>.-- The 50-hour electron capturing americium activity was tentatively assigned<sup>1</sup> to Am<sup>240</sup>. This assignment has been confirmed. The 50-hour activity is produced in good yield from the bombardment of Pu<sup>239</sup> with 10-Mev deuterons and is not observed in the bombardment of Pu<sup>239</sup> with 9-Mev protons (60-inch cyclotron) in which Am<sup>239</sup> is produced in good yield. Since the yield of the (d,3n) reaction is low at 10 Mev and since (p,n) should be the principle reaction producing americium with 9-Mev protons, the 50-hour activity is best assigned to Am<sup>240</sup>. No alpha-particles associated with the 50-hour americium have been observed. This is consistent with its assignment to Am<sup>240</sup> since the systematics of alpha-radioactivity<sup>4</sup> predict for this odd-odd nucleus a partial half-life for alpha-decay of some  $10^3$  years.

Am<sup>241</sup>.-- The production of this isotope in milligram amounts by the neutron irradiation of plutonium<sup>5</sup> makes it possible to study the higher mass isotopes of americium produced by (n, $\gamma$ ) reactions and the results of such experiments are reported in the following paragraphs. Recent specific activity measurements by B. B. Cunningham, S. G. Thompson, and H. R. Lohr<sup>6</sup> give a somewhat shorter

---

<sup>4</sup>I. Perlman, A. Ghiorso, and G. T. Seaborg, Phys. Rev. 77, 26 (1950).

<sup>5</sup>A. Ghiorso, R. A. James, L. O. Morgan, and G. T. Seaborg, Phys. Rev., in press.

<sup>6</sup>B. B. Cunningham, S. G. Thompson, and H. R. Lohr, unpublished work.



half-life for  $\text{Am}^{241}$  than that previously reported.<sup>7</sup> The new half-life is 475 years.

$\text{Am}^{242}$ .-- In addition to the 16-hour  $\text{Am}^{242m}$ , a long-lived  $\text{Am}^{242}$  produced by the  $(n, \gamma)$  reaction on  $\text{Am}^{241}$  has been reported;<sup>1</sup> alpha-decay of this isotope was demonstrated by separating the  $\text{Np}^{238}$  daughter and 0.5-Mev beta-particles in the americium fraction were also ascribed to  $\text{Am}^{242}$ . The fact that the observed beta-particles do belong to  $\text{Am}^{242}$  has now been demonstrated by observing the growth of  $\text{Cm}^{242}$  (half-life 162 days) in the purified, pile neutron-bombarded americium in an amount corresponding to the number of beta-particles observed. Mass spectrographic analysis of this americium shows  $\text{Am}^{242}$  to be present to the extent of ca. 0.5 percent. This analysis together with the results from the above  $\text{Cm}^{242}$  growth experiment and the yield from a  $\text{Np}^{238}$  extraction allow both the alpha- and beta-decay half-lives of  $\text{Am}^{242}$  to be estimated. The partial half-life obtained for beta-particle emission is roughly  $10^2$  years and that for alpha-particle emission is ca.  $10^4$  years. The  $(n, \gamma)$  cross section for the formation of  $\text{Am}^{242}$  is very roughly  $10^{-22}$   $\text{cm}^2$  but this may be off by a factor of several due to the many uncertainties involved, especially the neutron flux.

$\text{Am}^{243}$ .-- Neptunium separations from the americium fraction of an irradiation of  $\text{Am}^{241}$  with pile neutrons show the presence of equilibrium amounts of both  $\text{Np}^{238}$  and  $\text{Np}^{239}$ . The presence of  $\text{Np}^{239}$  proves the existence of the new isotope  $\text{Am}^{243}$  which is produced by two successive neutron capture processes in  $\text{Am}^{241}$  and decays by alpha-particle emission to  $\text{Np}^{239}$ . Mass spectrographic analysis of the americium of this bombardment showed  $\text{Am}^{243}$  present to the extent of ca. 0.5 percent. This together with the yield of  $\text{Np}^{239}$  determined in the chemical

---

<sup>7</sup>B. B. Cunningham, National Nuclear Energy Series, Plutonium Project Record, Vol. 14B, "The Transuranium Elements: Research Papers," Paper No. 19.2 (McGraw-Hill Book Co., Inc., New York, 1949).

extraction experiments gives a partial half-life for alpha-particle emission for  $\text{Am}^{243}$  of roughly  $10^4$  years. Alpha-particle pulse analysis of americium from another irradiation (containing approximately 10 percent  $\text{Am}^{243}$ ) showed the alpha-particle energy to be  $5.21 \pm 0.03$  Mev. This energy and half-life indicate that alpha-emission in this odd-even nucleus is not prohibited,<sup>4</sup> but like the case of the analogous nucleus  $\text{Am}^{241}$ , the measured alpha-particle energy may not represent the ground state transition. If one takes  $5.84$  Mev<sup>8</sup> for the alpha-particle energy of the ground state transition of  $\text{Cm}^{243}$  and closes a decay cycle using the measured alpha-particle energy of  $\text{Am}^{243}$  and  $0.7$  Mev for the beta-decay disintegration energy<sup>9</sup> of  $\text{Np}^{239}$ , one finds that  $\text{Am}^{243}$  could be unstable with respect to beta-decay by ca.  $0.05$  Mev. However, due to the uncertainties in the above energies (especially  $\text{Np}^{239}$ ) it is entirely possible that  $\text{Am}^{243}$  is beta-stable; the alpha-particle energy of  $\text{Cm}^{243}$  is also very uncertain and it is not known whether it corresponds to the ground state transition. In order to test this point curium was separated from an aged sample containing  $\text{Am}^{243}$  and no detectable  $\text{Cm}^{243}$  was found; this experiment places a lower limit of about  $10^3$  years on the partial half-life of  $\text{Am}^{243}$  for negative beta-emission. The cross section for the reaction  $\text{Am}^{242}(n,\gamma)\text{Am}^{243}$  seems to be large, of the order of some  $4 \times 10^{-21}$   $\text{cm}^2$ , a value which is subject to large error because of the uncertainty in the neutron flux.

$\text{Am}^{244}$ ? -- Irradiation of americium containing approximately 10 percent of the isotope  $\text{Am}^{243}$  with thermal neutrons in the uranium-heavy water pile at the Argonne National Laboratory produced a new americium activity of ca. 25-min. half-life at a yield corresponding to a cross section of roughly  $1/2 \times 10^{-22}$   $\text{cm}^2$ .

---

<sup>8</sup>K. Street, Jr., S. G. Thompson, and A. Ghiorso (to be published).

<sup>9</sup>G. T. Seaborg and I. Perlman, Revs. Mod. Phys. 20, 585 (1948).

This activity is probably due to the beta-emitting isotope  $\text{Am}^{244}$ , formed by an  $(n,\gamma)$  reaction.

We would like to thank Dr. R. A. James and Dr. S. G. Thompson for their assistance in many of the experiments, and Mr. F. L. Reynolds for the mass spectrographic analysis of the americium samples. It is a pleasure to thank T. M. Putnam, Jr., G. B. Rossi, J. T. Vale, and the crews of the 60-inch and 184-inch cyclotrons for their cooperation in the bombardments with these machines, and also W. H. Zinn and the operating group of the Argonne Laboratory pile for their cooperation and help in neutron irradiation experiments.

This work was performed under the auspices of the United States Atomic Energy Commission.